


What we need to know about NGS in Hi-C

François Serra, David Castillo & Marc A. Marti-Renom
Structural Genomics Group (CNAG-CRG)




Hi-C experiment




HiC mapping



Iterative
mapping




HiC mapping



HiC mapping




Iterative mapping




HiC mapping




Iterative
mapping




HiC mapping




HiC mapping




HiC mapping




HiC mapping




HiC mapping



HiC mapping





HiC mapping



How much do we map?

- 80-90% each end => 64-81% intersection
- 1% multiple contacts
- many of these will be lost in the filtering...



Dynabeads with streptavidin

TTGCAA

Restriction enzyme (RE) site

TTGCATCGAA



RE ligation site, repaired nucleotides in yellow (new cytosins are byotiniated)





Genomic single strand regions





read fragments from Hi-C



- Dynabeads with streptavidin
- Restriction enzyme (RE) site
- RE ligation site, repaired nucleotides in yellow (new cytosins are biotinylated)
- Genomic single strand regions
- read fragments from Hi-C



-  Dynabeads with streptavidin
-  **TTGCAA** Restriction enzyme (RE) site
-  **TTGCATCGAA** RE ligation site, repaired nucleotides in yellow (new cytosins are byotiniated)
-  Genomic single strand regions
-  read fragments from Hi-C



-  Dynabeads with streptavidin
-  Restriction enzyme (RE) site
-  RE ligation site, repaired nucleotides in yellow (new cytosins are biotinylated)
-  Genomic single strand regions
-  read fragments from Hi-C

