

ADVANCED BIostatISTICS

ABSTAT18

EM Algorithm

Lisete Sousa

Department of Statistics and Operations Research | CEAUL
Faculty of Sciences of Lisbon University



IGC, April 3rd - 6th, 2018

Contents

Introduction

MLE - Allele frequencies from genotype frequencies

EM Algorithm - Allele frequencies from phenotype frequencies

EM Algorithm - Differentially expressed genes

ESTIMATION OF ALLELE FREQUENCIES

... From Genotype Frequencies

- ▶ Maximum likelihood
- ▶ Example – Blood type
- ▶ Example in R

... From Phenotype Frequencies

- ▶ Maximum likelihood
- ▶ EM algorithm
- ▶ Complicated system to solve...
- ▶ Example – Blood type
- ▶ Example in R

IDENTIFICATION OF DIFFERENTIALLY EXPRESSED GENES

- ▶ EM algorithm for mixture of distributions
- ▶ Example 1 – Normal-Uniform – package `nudge`
- ▶ Example 2 – Gamma-Gamma-Gamma

Introduction

The Expectation-Maximization (EM) algorithm is a **parameter estimation method which falls into the general framework of maximum likelihood estimation**, and is applied in cases where part of the data can be considered to be incomplete, or “hidden”.

It is essentially an **iterative optimization algorithm** which, at least under certain conditions, will converge to parameter values at a local maximum of the likelihood function.

The EM algorithm has become a popular tool in statistical estimation problems involving **incomplete data**, or in problems which can be posed in a similar form, such as **mixture estimation** or **censored data**.

Maximum likelihood estimation of allele frequencies from genotype frequencies

Consider the ABO blood group system (3 alleles). Genotype frequencies, under Hardy-Weinberg equilibrium, are as follows:

AA	A0	BB	B0	AB	00
p^2	$2pr$	q^2	$2qr$	$2pq$	r^2

NOTE: Here, allele frequencies p_A , p_B and p_0 are p , q and r , respectively.

A set of n individuals contains:

- ▶ X_{AA} - Number of individuals with genotype AA;
- ▶ X_{A0} - Number of individuals with genotype A0;
- ▶ X_{BB} - Number of individuals with genotype BB;
- ▶ X_{B0} - Number of individuals with genotype B0;
- ▶ X_{AB} - Number of individuals with genotype AB;
- ▶ X_{00} - Number of individuals with genotype 00.

Thus,

$$(X_{AA}, X_{A0}, X_{BB}, X_{B0}, X_{AB}, X_{00}) \sim \text{Multinomial}(n; p^2, 2pr, q^2, 2qr, 2pq, r^2)$$

The corresponding likelihood function is

$$\begin{aligned}
 & L(p, q, r | n_{AA}, n_{A0}, n_{BB}, n_{B0}, n_{AB}, n_{00}) = \\
 & = \frac{n!}{n_{AA}! n_{A0}! n_{BB}! n_{B0}! n_{AB}! n_{00}!} \times \\
 & \quad \times (p^2)^{n_{AA}} (2pr)^{n_{A0}} (q^2)^{n_{BB}} (2qr)^{n_{B0}} (2pq)^{n_{AB}} (r^2)^{n_{00}} \propto \\
 & \propto p^{2n_{AA} + n_{A0} + n_{AB}} q^{2n_{BB} + n_{B0} + n_{AB}} r^{n_{A0} + n_{B0} + 2n_{00}}
 \end{aligned}$$

Note: $n = n_{AA} + n_{A0} + n_{BB} + n_{B0} + n_{AB} + n_{00}$

By Lagrange multiplier, we obtain the maximum likelihood estimates:

$$\blacktriangleright \hat{p} = \frac{1}{2n}(2n_{AA} + n_{A0} + n_{AB}) = (n_{AA} + \frac{1}{2}n_{A0} + \frac{1}{2}n_{AB}) / n$$

$$\blacktriangleright \hat{q} = \frac{1}{2n}(2n_{BB} + n_{B0} + n_{AB}) = (n_{BB} + \frac{1}{2}n_{B0} + \frac{1}{2}n_{AB}) / n$$

$$\blacktriangleright \hat{r} = \frac{1}{2n}(2n_{00} + n_{A0} + n_{B0}) = (n_{00} + \frac{1}{2}n_{A0} + \frac{1}{2}n_{B0}) / n$$

Example 1: For the data

AA	AB	A0	BB	B0	00
12	35	21	15	30	9

the maximum likelihood estimates are: $\hat{p} = 0.328$, $\hat{q} = 0.389$ and $\hat{r} = 0.283$.

How to do this in R?

Read the data:

```
> nAA<-12
> nAB<-35
> nA0<-21
> nBB<-15
> nB0<-30
> n00<-9
> n<-nAA+nAB+nA0+nBB+nB0+n00
```

Calculate de estimates:

```
> p<-(nAA+nA0/2+nAB/2)/n
> q<-(nBB+nB0/2+nAB/2)/n
> r<-(n00+nA0/2+nB0/2)/n
> p;q;r
```

```
[1] 0.3278689
```

```
[1] 0.3893443
```

```
[1] 0.2827869
```



EM algorithm for estimating the allele frequencies from phenotype frequencies

Situation: Grouped Data


The estimation is relatively easy if the genotype frequencies are known. However, this is not usual. For ABO blood group system, for example, what we frequently know is the blood type.

Example 1 (cont.):

The previous data would then be:

A	B	AB	0
33	45	35	9

The variables representing the number of individuals with each genotype and phenotype, can be organized according to the following scheme:



How to estimate p , q and r in this case?

Consider the model:

A	B	AB	0
$p^2 + 2pr$	$q^2 + 2qr$	$2pq$	r^2

Maximum likelihood estimation:

A complicated equation system.

EM algorithm:

Useful when information is missing. In this particular case, there is no information about n_{AA} , n_{A0} , n_{BB} and n_{B0} .

Step 1: Choose initial values for p , q and r . Let them be represented by $p_{(0)}$, $q_{(0)}$ and $r_{(0)}$.

Step 2 (E): Calculate an approximate value for the missing terms, using the previous initial values.

$$\blacktriangleright n_{AA}^* = n_A \frac{p_{(0)}^2}{p_{(0)}^2 + 2p_{(0)}r_{(0)}} = n_A \frac{p_{(0)}}{p_{(0)} + 2r_{(0)}}$$

Note that $X_A \sim \text{Bi}(n, p^2 + 2pr)$ and $X_{AA} | X_A \sim \text{Bi}(n_A, \frac{p^2}{p^2 + 2pr})$.

$$\blacktriangleright n_{A0}^* = n_A \frac{2r_{(0)}}{p_{(0)} + 2r_{(0)}}$$

$$\blacktriangleright n_{BB}^* = n_B \frac{q_{(0)}}{q_{(0)} + 2r_{(0)}}$$

$$\blacktriangleright n_{B0}^* = n_B \frac{2r_{(0)}}{q_{(0)} + 2r_{(0)}}$$

Step 3 (M): Calculate the first iterate for p , q and r , from maximum likelihood estimators. Here, missing information is substituted by the approximate values obtained in the previous step.

$$\blacktriangleright p_{(1)} = \frac{1}{2n}(2n_{AA}^* + n_{A0}^* + n_{AB})$$

$$\blacktriangleright q_{(1)} = \frac{1}{2n}(2n_{BB}^* + n_{B0}^* + n_{AB})$$

$$\blacktriangleright r_{(1)} = \frac{1}{2n}(2n_{00} + n_{B0}^* + n_{A0}^*)$$

Step 4: Repeat step 2, substituting $p_{(0)}$, $q_{(0)}$ and $r_{(0)}$ by $p_{(1)}$, $q_{(1)}$ and $r_{(1)}$.

Step 5: Repeat the procedure until convergence.

Initial values:

1. Start with 0.3(3) for $p_{(0)}$, $q_{(0)}$ and $r_{(0)}$.
2. Bernstein's suggestion (1925):

$$p_{(0)} = 1 - \sqrt{(n_0 + n_B)/n}$$

$$q_{(0)} = 1 - \sqrt{(n_0 + n_A)/n}$$

$$r_{(0)} = \sqrt{n_0/n}$$

Example 1 (cont.):

EM algorithm estimates are $\hat{p} = 0.331$, $\hat{q} = 0.409$, $\hat{r} = 0.259$.

How to do this in R?

Read the data:

```
> nA<-33  
> nB<-45  
> nAB<-35  
> n0<-9
```

Initialize the parameters (Bernstein's suggestion) – Step 1:

```
> p<-1-sqrt((n0+nB)/n)  
> q<-1-sqrt((n0+nA)/n)  
> r<-sqrt(n0/n)  
> p;q;r
```

```
[1] 0.3347009  
[1] 0.4132613  
[1] 0.2716072
```


Calculate approximate values for n_{AA} , n_{A0} , n_{BB} , n_{B0} – Step 2 (E):

```
> exp<-function(p,q,r){  
+ nAA<-nA*p/(p+2*r)  
+ nA0<-nA*2*r/(p+2*r)  
+ nBB<-nB*q/(q+2*r)  
+ nB0<-nB*2*r/(q+2*r)  
+ c(nAA,nA0,nBB,nB0)  
+ }
```

Visualize n_{AA}^* , n_{A0}^* , n_{BB}^* , n_{B0}^* , obtained in the first iteration:

```
> exp(p,q,r)  
[1] 12.58109 20.41891 19.44300 25.55700
```

NOTE that the first two values sum to 33 (n_A) and the last two sum to 45 (n_B).

Update p , q and r – Step 3 (M):

```
> param<-function(nAA,nA0,nBB,nB0){  
+ p<-(2*nAA+nA0+nAB)/(2*n)  
+ q<-(2*nBB+nB0+nAB)/(2*n)  
+ r<-(2*n0+nA0+nB0)/(2*n)  
+ c(p,q,r)  
+ }
```

Visualize \hat{p} , \hat{q} and \hat{r} , obtained in the first iteration:

```
> param(nAA,nA0,nBB,nB0)  
[1] 0.3278689 0.3893443 0.2827869
```

Iterative procedure – Steps 4 and 5:

```
> i<-0; er<-1
> while(sum(er>=0.00001)>0)
+ { e<-exp(p,q,r) # Step E
+   par<-param(e[1],e[2],e[3],e[4]) # Step M
+   er<-abs(c(p,q,r)-c(par[1],par[2],par[3])) # STOP criteria
+   i<-i+1
+   p<-par[1]; q<-par[2]; r<-par[3]
+   cat(i,p,q,r,"\n") # print the 3 values for each iteration.
+ }
```

```
1 0.3302504 0.4075533 0.2621964
2 0.3309501 0.4085211 0.2605288
3 0.3312228 0.4089185 0.2598587
4 0.3313321 0.4090797 0.2595882
5 0.3313762 0.4091449 0.2594789
6 0.331394 0.4091713 0.2594346
7 0.3314012 0.409182 0.2594167
8 0.3314042 0.4091863 0.2594095
```

Print the final solution in a friendly way:

```
> cat("\n Obtained solution after ",i," iterations:\n  
+ p* =",p,  
+ "\n q* =",q,  
+ "\n r* =",r,"\n")
```

Obtained solution after 8 iterations:


```
p* = 0.3314042  
q* = 0.4091863  
r* = 0.2594095
```

In example 1, the estimates of the allele frequencies obtained from both methods are very similar:

	MLE (known genotypes)	EM (unknown genotypes)
p^*	0.328	0.331
q^*	0.389	0.409
r^*	0.283	0.259

Final Remarks

For the situation in which we have grouped data, the EM algorithm may be summarized according to the following scheme:



EM algorithm on the identification of differentially expressed genes

Situation: Mixture of distributions

EM algorithm can be applied to other situations, which require more probabilistic calculations. That is the case when we have a mixture of distributions. As these calculations require a more advanced level of theoretical knowledge, we will just show the main idea through some publications and examples:

Normal – Uniform Distributions

BMC Bioinformatics



Methodology article

Open Access

Normal uniform mixture differential gene expression detection for cDNA microarrays

Nema Dean* and Adrian E Raftery

Address: Department of Statistics, University of Washington, Box 354322, Seattle, WA 98195-4322, U.S.A

Email: Nema Dean* - nemad@stat.washington.edu; Adrian E Raftery - raftery@stat.washington.edu

* Corresponding author

Published: 12 July 2005

Received: 17 March 2005

BMC Bioinformatics 2005, 6:173 doi:10.1186/1471-2105-6-173

Accepted: 12 July 2005

This article is available from: <http://www.biomedcentral.com/1471-2105/6/173>

© 2005 Dean and Raftery; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

- ▶ Dean and Raftery (2005), consider a **Normal Uniform mixture model** (NUDGE - Normal Uniform Differential Gene Expression) for the normalized log intensities.
- ▶ The method uses **EM algorithm** to estimate membership probabilities for each gene and classify it accordingly.
- ▶ The mixture gives **posterior probabilities of differential expression** which do not need to be adjusted for multiple testing.
- ▶ Typically a **threshold for probability** of being expressed of 0.5 is used for classification of the genes into the two groups (prob. ≥ 0.5 for DE genes; prob. < 0.5 for non DE genes).

Gamma – Gamma – Gamma Distributions

824

Biometrical Journal 50 (2008) 5, 824–836 DOI: 10.1002/bimj.200710468

Bayesian Classification and Non-Bayesian Label Estimation via EM Algorithm to Identify Differentially Expressed Genes: a Comparative Study

Marília Antunes^{*} and Lisete Sousa

University of Lisbon, Faculty of Sciences and Center of Statistics and Applications, DEIO, C6, Piso 4,
1749-016 Lisboa, Portugal

Received 31 October 2007, revised 7 March 2008, accepted 30 May 2008

- ▶ Antunes and Sousa (2008) consider a simple mixture model at the data level, as in Dean and Raftery (2005), with the difference that they consider that each gene is either down-regulated, non-DE or up-regulated and **three Gamma distributions** are assumed for the intensity measures.
- ▶ In addition to the **EM algorithm**, the authors apply a second method for gene classification, which is based on an hierarchical Bayesian model.

Acknowledgements:

Antónia Turkman (DEIO-FCUL), for allowing the use of some material produced by us in previous courses.

Bibliography:

Antunes, M. and Sousa, L. (2008). Bayesian classification and non-Bayesian label estimation via EM algorithm to identify differentially expressed genes: a comparative study. *Biometrical Journal* 50(5), 824-836.

Dean, N. (2006). The Normal Uniform Differential Gene Expression: (nudge) Detection Package. Bioconductor.

Dean, N. and Raftery, A.E. (2005). Normal uniform mixture differential gene expression detection for cDNA microarrays. *BMC Bioinformatics* 6, 173.

McLachlan, G. and Krishnan, T. (2008). *The EM Algorithm and Extensions*, 2nd Edition, Wiley.

Smyth, G.K., Ritchie, M., Thorne, N., Wettenhall, J. and Shi, W. (2016). User's Guide. limma: Linear Models for Microarray Data. Bioconductor.