Advanced Biostatists - ABSTAT18

IGC, 3rd - 6th April, 2018

PROJECT - PCA and Bootstrap.

The dataset "meaudret" can be found in package ade4. It is a classical set of ecological information: three data frames containing environmental variables, Trichopters abundance collected on 5 sites during 4 seasons along the Meaudret river (a river near Grenoble, France). The data set contains physico-chemical variables measured on 5 sites along the Meaudret River during four seasons (Pegaz-Maucet, 1980). Water samples were taken to study the following physico-chemical variables:

- Temp Water Temperature (in Celcius degree)
- Flow Flow (in l/s)
- pH pH
- Cond Conductivity (in μ S/cm)
- Bdo5 Biological oxygen demand after days (in mg/l)
- Oxyd Oxygen (in mg/l of oxygen)
- Ammo Ammonium Hydroxide (in mg/l NH_4^+)
- Nitr Nitrate (in mg/l NO_3^-)
- Phos Phosphate (in mg/l PO_4^{---})

```
library(ade4)
data(meaudret)
names(meaudret)
[1] "env" "design" "spe" "spe.names"
names(meaudret$mil)
[1] "Temp" "Flow" "pH" "Cond" "Bdo5" "Oxyd" "Ammo" "Nitr" "Phos"
```

meaudret\$env is a data frame containing all the environmental variables (9 variables in this case). meaudret\$design is a data frame detailing the experimental design. meaudret\$spe is a data frame providing information on species richness per site, per season.

There are 4 measures (one per season) taken per site and 5 measures (one per site) for each season.

- 1. Use a Non-parametric Bootstrap approach for the following:
 - (a) Estimate the median for the pollution variable (Ammo) using a 99% CI.
 - (b) Analyze if there are significant differences between pH mean values of the water in the winter and autumn?
- 2. What findings can you get from the data using PCA analysis (e.g. are the pollution variables more related with a specific season, or site).

Additional information

- You will have 10' to make your presentation.
- After your presentation there will be a discussion of your results.

References

D. Pegaz-Maucet. Impact d'une perturbation d'origine organique sur la d'erive des macro-invertebres benthiques d'un cours d'eau. Comparaison avec le benthos. PhD thesis, 1980.