



GTPB

The Gulbenkian Training Programme in Bioinformatics
(Since 1999)

Pedro Fernandes, Organiser



ELB19₉F

Entry Level Bioinformatics

04-08 February 2019

(First 2019 run of this Course)

Basic Bioinformatics Sessions

Practical 1: Databases and Tools

Wednesday 30 January 2019

Investigating the gene(s) associated with Aniridia

As a starting point for this exercise, imagine you have a vital interest in discovering and investigating the main human gene responsible for the terrible disease of the eye, **Aniridia**. There are many ways (including **google!**) you could discover what this gene might be. I choose to delve into the vast seas of knowledge so generously proffered by the **The National Center for Biotechnology Information (NCBI)**.


So, begin by going to the **Home Page** of the **The National Center for Biotechnology Information (NCBI)** (“<http://www.ncbi.nlm.nih.gov/>”).

You will arrive at a page offering access to the many **NCBI** resources available to you. Currently, you only require to search for genes, specifically those that relate to **Aniridia**, so first set the database selection field of the **Search** facility at the top of your page to **Gene**, set the **Search** field to **Aniridia** and click on the **Search** button.



A fine list of genes will emerge, including those sought. However, our interest is specific to **Human**, so the search should really be organism specific. To do this, one needs to execute an **Advanced** search. So, click on the **Advanced** button of the **Search** tool.

Now you can specify the precise field(s) of the annotation you wish to interrogate. In this case, set the **Disease/Phenotype** field to **Aniridia** and the **Organism** field to **Human**. As the two conditions are linked by **AND**, both must be true for any gene to be listed.



Click on the pretty red **Search** button.

Name/Gene ID	Description	Location	Aliases	MIM
<input type="checkbox"/> WT1 ID: 7490	Wilms tumor 1 [<i>Homo sapiens</i> (human)]	Chromosome 11, NC_000011.10 (32387775..32435539, complement)	AWT1, GUD, NPFS4, WAGR, WIT-2, WT33	607102
<input type="checkbox"/> PAX6 ID: 5080	paired box 6 [<i>Homo sapiens</i> (human)]	Chromosome 11, NC_000011.10 (31784792..31817961, complement)	AN, AN2, ASGD5, D11S812E, FVH1, MGDA, WAGR	607108
<input type="checkbox"/> TRIM44 ID: 54765	tripartite motif containing 44 [<i>Homo sapiens</i> (human)]	Chromosome 11, NC_000011.10 (35662692..35811053)	AN3, DIPB, HSA249128, MC7	612298
<input type="checkbox"/> ELP4 ID: 26610	elongator acetyltransferase complex subunit 4 [<i>Homo sapiens</i> (human)]	Chromosome 11, NC_000011.10 (31509729..31784525)	AN, AN2, C11orf19, PAX6NEB, PAXNEB, dJ68P15A.1, hELP4	606985
<input type="checkbox"/> DEL11P13 ID: 100528024	Wilms tumor, aniridia, genitourinary anomalies and mental retardation syndrome [<i>Homo sapiens</i> (human)]		C11DELP13, WAGR	194072

Just a few genes survive. All should really be examined, but this is just an exercise, so trust me ... it is **PAX6** that is the most interesting gene¹, in this context. This is the one to follow up by clicking on the link to its details.

From the **Summary** section one can conclude (sticking to the features that pertain to this exercise) that:

- there are two major domains, a paired domain and a homeobox, both of which bind DNA
- the gene regulates transcription (is a transcription factor)
- there is more than one protein isoform, and thus more than one transcript variant.

Summary This gene encodes a homeobox and paired domain-containing protein that binds DNA and functions as a regulator of transcription. Activity of this protein is key in the development of neural tissues, particularly the eye. This gene is regulated by multiple enhancers located up to hundreds of kilobases distant from this locus. Mutations in this gene or in the enhancer regions can cause ocular disorders such as aniridia and Peter's anomaly. Use of alternate promoters and alternative splicing result in multiple transcript variants encoding different isoforms. [provided by RefSeq, Jul 2015]

¹ This despite **WT1** being at the top of the list? This is a new promotion for **WT1**. For years it has been but a close second to **PAX6**. Whilst congratulations are clearly in order, this elevation is jolly inconvenient for the story I wish to reveal. So ... I intend to ignore it!

From the **Genomic context** section it can be seen that:

- **PAX6** is situated on **Chromosome 11**, band **p13**
- **PAX6** is on the complementary strand relative to that chosen to represent **Chromosome 11**
- **ELP4** (another human gene listed as associated with **Aniridia**) is very close, on the opposite strand to **PAX6**. This might be worthy of a mention, at a later time?
- There are **17** exons for **PAX6**. Jolly good, but I really wanted to know how many transcripts there were according to the **NCBI**? That is, how many different ways it is thought that nature spliced the **17** exons together. I would also like to discover how many distinct **isoforms** the **NCBI** imagines to result from however many **transcripts**. I proceed with impatience!

See PAX6 in [Genome Data Viewer](#)

Location: 11p13
Exon count: 17

Annotation release	Status	Assembly	Chr	Location
109	current	GRCh38.p12 (GCF_000001405.38)	11	NC_000011.10 (31784792..31817961, complement)
105	previous assembly	GRCh37.p13 (GCF_000001405.25)	11	NC_000011.9 (31806340..31839509, complement)

Click the **Genome Data Viewer** link. The **PAX6** genomic region, as interpreted by the **NCBI Genome Database**, is displayed for your delectation.

So, if I tell you the region displayed is the entire **PAX6** region of **Chromosome 11** and the green lines labelled on the right as something beginning with **NM_** represent the different transcripts, **can you now say how many transcripts there are according to this view?** In passing, the blobs along each line represent the exons. Dark blobs are coding exons. Light blobs represent the exons that form the **3'/5' UTR** regions of each transcript. The Introns are the pale green lines joining the blobs together.

The prediction of the transcripts shown here are based on database searches of all Human mRNA sequences stored in **RefSeq** against this region of the genome. The theory is that every Human mRNA sequence must match (nearly) perfectly somewhere in the human genome. Where it matches, there must be the genomic DNA from which the mRNA was transcribed. How charmingly true and simple!

To differentiate between coding and non-coding exons of a transcript, why not compare all human proteins with the genome (after suitable translation to amino acid codes in all six reading frames). They too must match near perfectly somewhere, identifying the **CoDing Sequence (CDS)** of each transcript. Transcript fully located. Job done! Of course, it does not always work so very neatly, but we need not admit that for the moment at least.

Comparing proteins with the genome is clumsy, compute intensive, slow. For major organisms (currently just **Human** and **Mouse**), specially compiled comprehensive databases of extremely reliable **DNA Coding Sequences** have been constructed. Searching with these databases enables much more efficient searching for coding exons and so is very much preferred.

OK, times up, the correct number of **PAX6** transcripts, according to the evidence you are offered here is **11**, of course! A conclusion you came to by counting the transcript prediction lines in the display. Jolly good! But it should be noted that the transcript count (and much else) depends on many transient circumstances, including particularly the versions of the databases used to build the views.

Quite recently indeed, **RefSeq** included **9** extra **PAX6** mRNA sequences of less certainty than the ones you see represented above. As more evidence was gathered, the credibility of these “extra” mRNA sequences was proved insufficient and they were removed. However, while they existed, they increased the transcript count to **20**!


This exposes that many of the “*facts*” presented by these services are but “*predictions*” that will vary as more/better data become available. Pretty good predictions, but nevertheless, *predictions*!

As will be emphasised throughout these exercises, databases in general contain entries (often simply predictions) of varying certainty. That being so, the user must be able to ascertain the relative quality of a given data item. In the case of mRNA sequences in **RefSeq**, the entry **Accession Codes** (unique data entry labels) indicate the quality of the evidence for the mRNA predictions. **Accession Codes** which begin with **XM_** indicate mRNA sequence predictions of less certainty than those that begin **NM_**. Typically, the less certain entries (**XM_** entries) have been identified by computer programs alone. The **NM_** entries, normally, will have been properly investigated by human experimenters/investigators.

It gets worse! Other factors interfere with any hope of simple answers to seemingly trivial inquiries such as “how many transcripts are there?”. One such factor being *where* the question is asked.

Move back to the page describing the **PAX6** gene. In the familiar graphic at the top of the **Genome regions, transcripts and products** section you will find routes to corresponding information from the **Ensembl Genome Database**. Hover over the **PAX6** (also known as **ESNG00000007372**, by **Ensembl** and close friends) green line in the bottom half of the picture. You will be rewarded by cheery grey box full of links to **Ensembl** and other exciting places.

This gene has 84 transcripts ([splice variants](#)), [145 orthologues](#), [50 paralogues](#), is a member of [2 Ensembl protein families](#) and is associated with [30 phenotypes](#).




Use the link labelled **View ENSEMBL**:. A view of the region of **Chromosome 11** similar to those you have already considered will leap forth. As before, the exons for each transcript are represented by blobs (filled for coding, empty for UTR regions). Introns are represented by wiggly lines joining the blobs. Notice first that there are considerably more than **11** transcripts represented here! At the top of the page, in tiny letters it claims **84**! (a massive increase even from the **31** transcripts predicted by a recent previous version of **Ensembl**!).

You *could* check this assertion by counting all the transcripts represented in the graphic, but I would not recommend doing so. Sometimes it is best just to believe. There are indeed **84**.

The colour scheme used for the transcripts [we might discuss in overview later](#). For now, just know that the **gold** transcripts are supported by better evidence than the **red** ones.

Once more a database that offers data items (“*predictions*”) of varying credibility.



Looking a little further down the transcripts displays, you will see that an increasing proportion of the transcripts are not **protein coding** (the blue ones). The display you examined at the **NCBI** only represented protein coding transcripts. This partially explains why **Ensembl** appears finds so many more transcripts that its broad alternatives.

So a further reason for not finding a consistent answer to the simple question “How many transcripts are there for the **PAX6** gene” is variation in the *definition* of a transcript.

Also, and more importantly, **Ensembl** and the **NCBI Genome Database** use different strategies to predict transcripts (and nearly everything else!). Both use database searches broadly in the manner described above and (for the human genome at least) the same basic assemblies of the genome and sequence databases. It is primarily the

interpretation of the data and analytical results that varies.

The database searches used as the fundamental strategy to identify transcripts take a very long time to execute, even given the immense computing resources available to the **NCBI** and the **Ensembl** teams. Some clever strategies are employed to minimise the time spent on these searches. It would be good to consider these, specifically with respect to their implementation by **Ensembl**, at least superficially.

For a more detailed view of the predicted transcripts, click on the **Show transcript table** link. The transcript predictions are now presented in the form of a table. The protein coding transcripts are all at the top of the table. I counted **56**, but I

Name	Transcript ID	bp	Protein	Biotype	CCDS	UniProt	RefSeq	Flags
PAX6-245	ENST00000638914.2	7290	422aa	Protein coding	CCDS31451	P26367 @ Q66SS1	-	GENCODE basic APPRIS ALT1
PAX6-270	ENST00000640368.1	6975	436aa	Protein coding	CCDS31452	F1T0F8 @ P26367	-	GENCODE basic APPRIS ALT1
PAX6-283	ENST00000643871.1	6944	422aa	Protein coding	CCDS31451	P26367 @ Q66SS1	NM_000280 @ NM_001258464 @ NP_000271 @ NP_001245393	GENCODE basic APPRIS ALT1
PAX6-231	ENST00000606377.6	6901	436aa	Protein coding	CCDS31452	F1T0F8 @ P26367	-	TSL:1 GENCODE basic APPRIS ALT1
PAX6-209	ENST00000419022.6	6888	436aa	Protein coding	CCDS31452	F1T0F8 @ P26367	NM_001604 @ NP_001585	TSL:1 GENCODE basic APPRIS P4
PAX6-203	ENST00000379109.7	3182	422aa	Protein coding	CCDS31451	P26367 @ Q66SS1	-	TSL:2 GENCODE basic APPRIS ALT1
PAX6-273	ENST00000640610.1	2730	422aa	Protein coding	CCDS31451	P26367 @ Q66SS1	-	GENCODE basic APPRIS ALT1
PAX6-284	ENST00000645710.1	2888	436aa	Protein coding	CCDS31452	F1T0F8 @ P26367	NM_001258462 @ NP_001245391	GENCODE basic APPRIS ALT1
PAX6-259	ENST00000639916.1	2622	422aa	Protein coding	CCDS31451	P26367 @ Q66SS1	NM_001258465 @ NP_001245394	GENCODE basic APPRIS ALT1
PAX6-243	ENST00000638903.1	2620	436aa	Protein coding	CCDS31452	F1T0F8 @ P26367	-	GENCODE basic APPRIS P4
PAX6-207	ENST00000379129.7	2614	436aa	Protein coding	CCDS31452	F1T0F8 @ P26367	-	TSL:5 GENCODE basic APPRIS ALT1
PAX6-202	ENST00000379107.7	2579	436aa	Protein coding	CCDS31452	F1T0F8 @ P26367	-	TSL:5 GENCODE basic APPRIS ALT1
PAX6-208	ENST00000379132.6	2576	422aa	Protein coding	CCDS31451	P26367 @ Q66SS1	-	TSL:5 GENCODE basic APPRIS ALT1
PAX6-282	ENST00000640975.1	2553	436aa	Protein coding	CCDS31452	F1T0F8 @ P26367	NM_001310158 @ NP_001297087	GENCODE basic APPRIS P4
PAX6-257	ENST00000639409.1	2450	436aa	Protein coding	CCDS31452	F1T0F8 @ P26367	NM_001258463 @ NP_001245392	GENCODE basic APPRIS P4

would not claim to be completely accurate, I wavered half way down the list! Lots more than the **NCBI** anyway.

Ensembl uses both the sequences of **RefSeq** mRNAs and those of their protein products (the **RefSeq** entries whose **Accession Codes** commence NP_) to predict transcripts, however, **Ensembl** appears to have less blind faith in the accuracy of these data than the **NCBI**.

Note: There is no “one to one” correspondence between **RefSeq** mRNAs and transcript predictions. All **11 RefSeq** mRNAs are referenced, but *two* are used to support the single third transcript in the list. If **Ensembl** regarded **RefSeq** mRNAs

as “perfect” (as the **NCBI** appears to do) this would clearly be nonsense! We should discuss why it is reasonable not to not to accept the infallibility of a **RefSeq** mRNA matches with the **Genome**.

PAX6-220	ENST00000525935.2	875	3aa	Protein coding	-	-	-	CDS 3' incomplete TSL:3
PAX6-260	ENST00000639920.1	676	72aa	Protein coding	-	A0A1W2PR58	-	CDS 3' incomplete
PAX6-256	ENST00000639394.1	1988	163aa	Nonsense mediated decay	-	A0A1W2PQW3	-	-
PAX6-227	ENST00000533156.2	848	No protein	Processed transcript	-	-	-	TSL:5
PAX6-213	ENST00000464174.6	846	No protein	Processed transcript	-	-	-	TSL:5
PAX6-222	ENST00000530373.6	785	No protein	Processed transcript	-	-	-	TSL:4
PAX6-223	ENST00000530714.6	650	No protein	Processed transcript	-	-	-	TSL:4
PAX6-267	ENST00000640251.1	649	No protein	Processed transcript	-	-	-	-
PAX6-229	ENST00000534353.5	540	No protein	Processed transcript	-	-	-	TSL:4
PAX6-254	ENST00000639203.1	532	No protein	Processed transcript	-	-	-	-
PAX6-233	ENST00000638278.1	417	No protein	Processed transcript	-	-	-	-
PAX6-275	ENST00000640617.1	412	No protein	Processed transcript	-	-	-	-
PAX6-279	ENST00000640819.1	368	No protein	Processed transcript	-	-	-	-
PAX6-228	ENST00000533333.5	6173	No protein	Retained intron	-	-	-	TSL:2
PAX6-216	ENST00000474783.2	4392	No protein	Retained intron	-	-	-	TSL:2
PAX6-214	ENST00000470027.7	3587	No protein	Retained intron	-	-	-	TSL:2
PAX6-265	ENST00000640172.1	2525	No protein	Retained intron	-	-	-	-

Looking further down the list you will see that many **Ensembl** protein coding transcripts are predicted without reference to any **RefSeq** entry.

Hover over the evidence **Flags** associated with the transcript predictions towards the end of the list. How reliable would you judge these predictions to be?

We could go on. Other sources (not necessarily **Genome Databases**) would count the transcripts differently again. Perhaps the best answer to the question “How many transcripts are there for the **PAX6** gene” is “**Several**”.

Before leaving **Ensembl**, it would be good to save the genomic sequence of this region for analysis later on.

To do this, first click on the **Sequence** link on the left hand side of the page. Under the transcript table the sequence of the **PAX6** region of the genome will be displayed. The exons will be tastefully highlighted for you delectation. The display includes **600** base pairs of **Flanking Sequence (3' and 5')** which are included (by default) when the sequence is downloaded

Marked-up sequence

Download sequence | BLAST this sequence

Exons PAX6 exons All exons in this region

Markup loaded

```
>chromosome:GRCh38:11:31784179:31818662:-1
ATACAATCACCTACATTTTCTAATGTGGTTGGAGCCTTTCAGCCAGAGGGCCGAGGGAAGC
CCGGGTAGGCCCTTTAGGGCTTCCCTCTTGAGAACCAGCAGGCTGGAGAGACCTTT
GGCCTAGGCCCTGAAAAAGGGGTCCGATGTCTCTTCCCGGAGCCCGCTGTGCCCCAG
CTAGTGACTTGGGGCTCGAGGGCCAGGTTGAGGGTACTCATCGAGCTCGAAGCTCTCC
TAAAAATGATTCTGCCAAAAGCGCTTCCATCCCAGGCGGGCTTCGGGTCTCTCCGA
TGAAAGGACTCCCTTGGGGATCGGAGGAGGGGACAGGGTGATTACCCAGAGAGGTAGCTG
GCCAGCCTAAGGGCAGAGATCTTGGGGCCCTAGTGCCGAAGGTGCGGAGGAGCGCACTC
GGCAAGACTAGTTTCTGGGGATCGACTCTACGCCATACAGGACGGGCGCCAGGCTGGA
CCGGCCGGGCTAGAGCAGTACAGGCGGGCCAAAGGCAAGGCAAGGAGGGTTGGAAGC
CGGCCGGACCTGGGTGGGGAGAAGCAGGCTCCCGCCGGCCGGAAGAACTAGTCGGCCGA
GAGCTGTGCCAACTTAGCCGCCATGACGTACGCGGGCCGGCCAGCCAATGAGGACGG
CGCTGGCGTGGATATTAAGGAAAGTTAGCGCCTGCTGAGCACCCCTTTTCTTATCATT
GACATTTAACTCTGGGGCAGGCTCTCGCTGAGAACCGGGCTGTCAGATCTGCCACTTCC
CCTGCCAGCGGGGTGAGAAGTGTGGAAACCAGGCTGCCAGGCTCACCTGCCCTCCCG
CCCTCCGCTCCCAAGTAAACCGCCGGGCTCCGGCCCGCCGGCTCGGGGCGCCGGGG
CCCTCCGCTGCCAGGACTGCTGTCCCAAAATCAAAGCCCGCCCAAGTGGCCCGGGG
CTTGATTTTGTCTTTAAAAGGAGGCATACAAAGATGGAAGCGAGTACTGAGGGAGGGA
TAGGAAGGGGGTGGAGGAGGGACTTGTCTTGGCCAGATGTGCTCTCTGCAAAAAGTAGC
AAAATGTTCCACTCCTAAGAGTGGACTTCCAGTCCGGCCCTGAGCTGGGAGTAGGGGGCG
GGAGTCTGCTGCTGCTGTCTGCTAAAGCCACTCGCCAGCCGCAAAAATGCAGGAGGTGGG
GACGCACTTTGCATCCAGACTCCTCTGCATCGCAGTTACGACATCCAGCTTGGGAAA
GTCCGTACCCCGCCCTGGAGCGCTTAAAGACACCTGCCCGGGTGGGGCAGGTTGCGACG
AGAAGTTTCCCGGGTGGCAAGTGCAGATGGCTGGACCGCAACAAAGTCTAGAGATGGG
GTTCTTTCTCAGAAAGACCGGAGTACGAAAGAAATGCCGGCCGACAGACTGGGAGCGC
GTAAGCTCCAGCGGTGATTGAGCTTCACTCGGAAGCAATAATTAGCGATTCT
```

File name:

File format:

Settings

Sequences to export:

- Select/deselect all
- cDNA (transcripts)
- Coding sequences (CDS)
- Amino acid sequences
- 5' UTRs
- 3' UTRs
- Exons
- Introns
- Genomic sequence

5' Flanking sequence (upstream): * (Maximum of 100000)

3' Flanking sequence (downstream): * (Maximum of 100000)

Now chose to **Download sequence**. The **Download sequence** form will burst into view.

Set the **File name:** to **pax6_genomic.fasta**

Set the **File format:** to **FASTA**

Accept the default **600** base pairs for both the **5' Flanking sequence (upstream):** and the **3' Flanking sequence (downstream):**.

Finally, click on the **Download** button and do whatever it takes to move the file you create to somewhere sensible on your **Desktop**.

Using whatever text editor is most convenient, edit your file to change the first word of the first line of the file to contain information, from **11** to **pax6_genomic**. This first word is defined as the sequence identifier in **FASTA** format (as, I hope, will be explained at some point). **pax6_genomic** is a far more informative identification than **11** (simply the Chromosome number).

```
>pax6_genomic| dna:chromosome chromosome:GRCh38:11:31784179:31818662:-1
ATACAATCACCTACATTTTCTAATGTGGTTGGAGCCTTTCAGCCAGAGGGCCGAGGGAAGC
CCGGGTAGGCCCTTTAGGGCTTCCCTCTTGAGAACCAGCAGGCTGGAGAGACCTTT
GGCCTAGGCCCTGAAAAAGGGGTCCGATGTCTCTTCCCGGAGCCCGCTGTGCCCCAG
CTAGTGACTTGGGGCTCGAGGGCCAGGTTGAGGGTACTCATCGAGCTCGAAGCTCTCC
TAAAAATGATTCTGCCAAAAGCGCTTCCATCCCAGGCGGGCTTCGGGTCTCTCCGA
TGAAAGGACTCCCTTGGGGATCGGAGGAGGGGACAGGGTGATTACCCAGAGAGGTAGCTG
GCCAGCCTAAGGGCAGAGATCTTGGGGCCCTAGTGCCGAAGGTGCGGAGGAGCGCACTC
GGCAAGACTAGTTTCTGGGGATCGACTCTACGCCATACAGGACGGGCGCCAGGCTGGA
CCGGCCGGGCTAGAGCAGTACAGGCGGGCCAAAGGCAAGGCAAGGAGGGTTGGGAC
CGCCGGACCTGGGTGGGAGAGGAGGCTCCCGCCGGCCGGCAAAACTAGTCTGGCGCA
GAGTGTGCCAACTTAGCCGCCATGACGTACGCGGGCCGGCCAGCCAATGAGGACGG
CGCTGGCGTGGATATTAAGGAAAGTTAGCGCCTGCTGAGCACCCCTTTTCTTATCATT
GACATTTAACTCTGGGGCAGGCTCCTCGGTAGAACCAGGGCTGTCAGATCTGCCACTTC
CCTGCCAGCGGGGTGAGAAGTGTGGAAACCAGGCTGCCAGGCTCACCTGCCCTCCCG
CCCTCCGCTCCAGGTAACCGCCGGGCTCCGGCCCGGGCCGGCTCGGGGCGCCGGGG
CCTCTCCGCTGCCAGGCTGCTGTCCCAAAATCAAAGCCCGCCCAAGTGGCCCGGGG
CTTGATTTTGTCTTTAAAAGGAGGCATACAAAGATGGAAGCGAGTACTGAGGGAGGGA
TAGGAAGGGGGTGGAGGAGGGACTTGTCTTGGCCAGTGTGCTCTGCAAAAAGTAGC
AAAATGTTCCACTCCTAAGAGTGGACTTCCAGTCCGGCCCTGAGCTGGGAGTAGGGGGG
GGAGTCTGCTGCTGCTGTCTGTAAGCCACTCGCAGCCCGGCAAAAATGAGGAGGTGGG
GACGCACTTTGCATCCAGACTCCTCTGCATCGCAGTTACGACATCCAGCTTGGGAAA
GTCCGTACCCCGGCTGGAGCGCTTAAAGACACCTGCCCGGGTGGGGCAGGTTGCGC
AGAAGTTTCCCGGGTGGCAAGTGCAGATGGCTGGACCGCAACAAAGTCTAGAGATGGG
```

The next investigation might be to discover “How many protein isoforms might there be for **PAX6**?”

Well, whilst the **Ensembl** transcript list is still in view, glance down the **Protein** column which displays the size of the protein products for each transcript. Clearly insufficient evidence for a serious **isoform** count, but enough to set a lower limit, as the same **isoform** cannot be more than one length! If there were not so very many! One might count how many different lengths of proteins were listed. I tried to do this, but I gave up around **twenty-something**. Let us be content to declare that there are **lots**. The most likely looking ones are either **422** or **436** amino acids long. Some of the others might cause a raised eyebrow or two, especially the one that is **3** amino acids long (third from last **Protein coding** entry in the list)? But, who are we to question! **Lots** is the informal **Ensembl** minimum total.

Click your way back to the **NCBI PAX6 gene entry**. So, now to discover the number of protein products (**isoforms**) that the **NCBI** predicts. This view makes this simple question clumsy to answer as the protein products of each transcript are reported separately (as they are by **Ensembl**), even when they are identical???

However, it can be done. Click on the **NCBI Reference Sequences (RefSeq)** link in the **Table of contents** on the right hand side of the page. Focus on the **mRNA and Protein(s)** sub-section. Skim down the entries for every transcript. Check the different isoform names. I see:

```
01 - NM_000280.4    → NP_000271.1    paired box protein Pax-6 isoform a
02 - NM_001127612.1 → NP_001121084.1 paired box protein Pax-6 isoform a
03 - NM_001258462.1 → NP_001245391.1 paired box protein Pax-6 isoform b
04 - NM_001258463.1 → NP_001245392.1 paired box protein Pax-6 isoform b
05 - NM_001258464.1 → NP_001245393.1 paired box protein Pax-6 isoform a
06 - NM_001258465.1 → NP_001245394.1 paired box protein Pax-6 isoform a
07 - NM_001310158.1 → NP_001297087.1 paired box protein Pax-6 isoform b
08 - NM_001310159.1 → NP_001297088.1 paired box protein Pax-6 isoform c
09 - NM_001310160.1 → NP_001297089.1 paired box protein Pax-6 isoform d
10 - NM_001310161.1 → NP_001297090.1 paired box protein Pax-6 isoform d
11 - NM_001604.5     → NP_001595.2    paired box protein Pax-6 isoform b
```

I count **4** different isoforms, imaginatively named **Isoform a**, **Isoform b**, **Isoform c** and **Isoform d**. One associated with each transcript description. Look carefully at the annotations and there is more information. In particular:

Description: **Isoform b** is also known as **Isoform 5a**. Why this is interesting will become apparent in a page or so.

Isoform b is also reported to be longer than **Isoform a**.

Description	Transcript Variant: This variant (5) differs in the 5' UTR and includes an alternate in-frame exon in the 5' coding region, compared to variant 1. The encoded isoform (b, also known as 5a) is longer than isoform a. Variants 2, 4, 5 and 8 encode the same isoform (b).
--------------------	--

Conserved Domains:

Conserved Domains (2) summary	
smart00351 Location:4 → 128	PAX; Paired Box domain
pfam00046 Location:214 → 266	Homeobox; Homeobox domain

Both **Isoform a** and **Isoform b** are recorded as having two domains. A **Paired Box Domain** at the beginning, and a **Homeobox Domain** further along.

Conserved Domains (2) summary	
smart00351 Location:4 → 142	PAX; Paired Box domain
pfam00046 Location:228 → 280	Homeobox; Homeobox domain

Both **Paired Box Domains** are primarily indicated by a hit with the relevant entry in the **SMART** database. Both **Homeobox Domains** are supported by matches with **Pfam** database entries. Other domain databases will almost certainly provide

supporting evidence, but reference to just one match is sufficient here.

From the location information, the **Paired Box** of **Isoform b** appears to include an extra **14** amino acids.

UniprotKB offers yet another version of this story. Just for a few clicks, let us intrude into the **UniProtKB** section of your course.

At the very bottom of the current page, you will find a link to **UniprotKB**. Use it.

Protein Accession	Links	
	GenPept Link	UniProtKB Link
P26367.2	GenPept	UniProtKB/Swiss-Prot:P26367.2

Lo! the **PAX6** human protein as seen and understood by **UniProtKB**. Click on the **Sequences (3+)** button on the left hand side of the page. **UniProtKB** declares **3** isoforms! At least, **3** that it is willing to admit to with certainty. Also mentioned are a further **33** that are suggested as possible by computer analysis.

Sequences (3+)
 Sequence status¹: Complete.
 This entry describes **3** isoforms¹ produced by **alternative splicing**. [Align](#) [Add to basket](#)
 This entry has 3 described isoforms and 33 potential isoforms that are computationally mapped. [Show all](#) [Align All](#)

There is **isoform 1**, also known as **isoform a** in America. Note that this is the “*canonical sequence*” for this protein. That is, this is the isoform used to represent this protein in **UniProtKB**. The sequence(s) of all other isoform(s) are recorded as elements of the annotation.

Isoform 1 (identifier: **P26367-1**) [[UniParc](#)] [FASTA](#) [Add to basket](#)
This isoform has been chosen as the 'canonical' sequence. All positional information in this entry refers to it. This is also the sequence that appears in the downloadable versions of the entry.

Also we have **Isoform 5a** (or **PAX6-5a**), also known as **isoform b** in America (where it also answers to **Isoform 5a** when pressed). Note that the entry declares the sequence difference to be:

47-47: Q → QTHADAKVQVLDNQN

Isoform 5a (identifier: **P26367-2**) [[UniParc](#)] [FASTA](#) [Add to basket](#)
 Also known as: Pax6-5a
 The sequence of this isoform differs from the canonical sequence as follows:
47-47: Q → QTHADAKVQVLDNQN

Literally:

“The amino acid at **position 47** is a **Q** in the canonical sequence. In **isoform 5a** this is replaced by the **15** amino acids **QTHADAKVQVLDNQN**”.

More coherently this amounts to:

“**isoform 5a** differs from the canonical **isoform 1** in that it has an insertion of **14** amino acids after the **47th** amino acid (a **Q**) of the canonical protein”.

It is significant to note that position **47** is right in the middle of the **Paired Box Domain** that occurs in both isoforms. This confirms that which was noticed at the **NCBI**.

Finally **UniProtKB** proudly presents the somewhat ephemeral **isoform 3** (or **PAX6-5A,6*** for those who enjoy formality). But, this one has no known sequence? Not much that Bioinformatics can offer here methinks.

Isoform 3 (identifier: **P26367-3**)
 Also known as: Pax6-5A,6*
 Sequence is not available

So I hope you will agree that the **UniProtKB** confident isoform count stands at a very modest **2**, plus a ghost.

To visualise the differences between the **2** isoforms with sequence, click on the **Align** button for the **3** described isoforms, at the top of the **Sequences** section. After deep thought and much fumbling, **UniProtKB** will multiply align all the selected isoform sequences for you. As there are only **2** in this case, this will appear very similar to a **Pairwise alignment**. Highlight the **DNA binding** regions and the **Domains**.

I leave the interpretation of this splendid display to you, and later short discussion if required.

The extra **14** amino acids of **isoform 5a** are due to the inclusion of a tiny extra (**42** base pair) exon in some transcripts.

Highlight

Annotation

- Alternative sequence
- Natural variant
- Domain**
- Sequence conflict
- DNA binding**
- Helix
- Compositional bias
- Turn
- Chain
- Beta strand

Alignment

[How to print an alignment in color](#)

P26367	PAX6_HUMAN	1	MQNSHSGVNQLGGVFVNGRPLPDSTROKIVELAHSGARPCDISRILQ-----	47
P26367-2	PAX6_HUMAN	1	MQNSHSGVNQLGGVFVNGRPLPDSTROKIVELAHSGARPCDISRILQTHADAKVQVLDNQ	60
P26367	PAX6_HUMAN	48	-VSNCGVSKILGRYYETGSIRPRAIGGSKPRVATPEVVSKIAQYKRECPSIFAWI	106
P26367-2	PAX6_HUMAN	61	NVSNCGVSKILGRYYETGSIRPRAIGGSKPRVATPEVVSKIAQYKRECPSIFAWI	120
P26367	PAX6_HUMAN	107	LSEGVCTNDNIPSVSSINRVLRLNASEKQMQMGADGMYDKLRMLNGTGSWGTRPGWY	166
P26367-2	PAX6_HUMAN	121	LSEGVCTNDNIPSVSSINRVLRLNASEKQMQMGADGMYDKLRMLNGTGSWGTRPGWY	180
P26367	PAX6_HUMAN	167	SVPGQPTQDGCQQQEGGENTNSISSNGEDSDEAQMRLQLKRKLQRNRTSFTQEQIEALE	226
P26367-2	PAX6_HUMAN	181	SVPGQPTQDGCQQQEGGENTNSISSNGEDSDEAQMRLQLKRKLQRNRTSFTQEQIEALE	240
P26367	PAX6_HUMAN	227	KEFERTHYPDVFAERLAALKIDLPEARIQVWFSNRRAKWRREEKLRNQRQASNTPSHIP	286
P26367-2	PAX6_HUMAN	241	KEFERTHYPDVFAERLAALKIDLPEARIQVWFSNRRAKWRREEKLRNQRQASNTPSHIP	300
P26367	PAX6_HUMAN	287	ISSSFSTSVYQPIPOPTTPVSSFTSGSMLGRDITALTNTYSALPPMPSFTMANNLPQPP	346
P26367-2	PAX6_HUMAN	301	ISSSFSTSVYQPIPOPTTPVSSFTSGSMLGRDITALTNTYSALPPMPSFTMANNLPQPP	360
P26367	PAX6_HUMAN	347	VPSQTSYSYSCMLPTSPSVNGRSYDITYTPPHMQTHMNSQPMGTSGLTISPGVSPVQ	406
P26367-2	PAX6_HUMAN	361	VPSQTSYSYSCMLPTSPSVNGRSYDITYTPPHMQTHMNSQPMGTSGLTISPGVSPVQ	420
P26367	PAX6_HUMAN	407	VPGSEPDMSQYWPRLLQ	422
P26367-2	PAX6_HUMAN	421	VPGSEPDMSQYWPRLLQ	436

Can you see the evidence for this assertion in the regional genomic maps of a few pages back?

We need to save a some protein sequences for future analysis. This is easiest from **UniProtKB** so now is good. To declare your intention to save the entire canonical version of the **PAX6** protein to a file, move back from your alignment. Move to the top of the page where you will find the bizarre invitation to **Add to basket** ? Just do it.

You also need to download the sequences of both domains is separate files, via your basket. First the **Paired Box**.

Click the **Family & Domains** button on the left of the page. Then use the **Add** button adjacent to the **Paired** entry. Its now in your basket you will be ecstatic to know.

Feature key	Position(s)	Description	Actions	Graphical view	Length
Domain ⁱ	4 - 130	Paired PROSITE-ProRule annotation	Add BLAST		127

As they are so conveniently in view, take note of the **Compositional bias** features. They will be of interest when we look at database searching.

Feature key	Position(s)	Description	Actions	Graphical view	Length
Compositional bias ⁱ	131 - 209	Gln/Gly-rich	Add BLAST		79
Compositional bias ⁱ	279 - 422	Pro/Ser/Thr-rich	Add BLAST		144

Natural variant ⁱ (VAR_008694)	29	I → S in AN.	1 Publication
Natural variant ⁱ (VAR_003811)	29	I → V in AN.	1 Publication
Natural variant ⁱ (VAR_008695)	33	A → P in AN.	1 Publication
Natural variant ⁱ (VAR_008696)	37 - 39	Missing in AN.	1 Publication
Natural variant ⁱ (VAR_008697)	42	I → S in AN; mild.	1 Publication
Natural variant ⁱ (VAR_008698)	43	S → P in AN.	1 Publication
Natural variant ⁱ (VAR_003812)	44	R → Q in AN.	1 Publication

Then take an excursion to glance at the **Pathology & Biotech** section. Note the many **Natural variants** recorded as responsible for AN (ANiridia, that is). Particularly those around amino acid positions **29** to **44** and specifically that at position **33**.

Looking at **PCR Primer Design** later, you will be attempting to create a **PCR** products from patients that, when sequenced, will determine the presence or absence of this variant.

Next, skip nimbly to the **Family & Domains** section. Concentrate on the **Family and domain databases** sub-section. Here are displayed the results of comparing the **PAX6** protein with many of the available **Domain/Motif Databases**, including those of the **Interpro Consortium**, collectively.

Are the results broadly as you might expect?

For an effective graphic summary, link to **View protein in InterPro** for the **Interpro** graphical results. If the detail is not entirely transparent, this result will be discussed further when you generate it for yourselves using **Interpro**.



The results you are looking at are computed, largely automatically, by the **UniProtKB/Interpro** annotation system. However, running many of the same analyses manually is trivial. Maybe you will do some in the course of these exercises?

Finally, return to the **UniProtKB PAX6** page and move to the **Structure** section.

Click on the **Show more details** button.

Describe the arrangement of Helices within **PAX6**.

Feature key	Position(s)	Description	Actions	Graphical view	Length
Beta strand ⁱ	6 - 8	Combined sources			3
Beta strand ⁱ	14 - 16	Combined sources			3
Helix ⁱ	23 - 34	Combined sources			12
Helix ⁱ	39 - 46	Combined sources			8
Helix ⁱ	50 - 63	Combined sources			14
Beta strand ⁱ	77 - 79	Combined sources			3
Helix ⁱ	81 - 93	Combined sources			13
Helix ⁱ	99 - 108	Combined sources			10
Turn ⁱ	114 - 116	Combined sources			3
Helix ⁱ	120 - 133	Combined sources			14
Helix ⁱ	219 - 229	Combined sources			11
Helix ⁱ	237 - 246	Combined sources			10
Helix ⁱ	251 - 275	Combined sources			25

Back to saving sequences for later! To get to the **Homeobox** domain, you need to click on the **Function** button on the left hand side of the page.

Feature key	Position(s)	Description	Actions	Graphical view	Length
DNA binding ⁱ	210 - 269	Homeobox PROSITE-ProRule annotation	Add BLAST		60

A valid question at this point might be “Why is the **Homeobox** domain a **Function** (specifically a **DNA binding** feature), but the **Paired** domain is a **Domain** feature?” To which the answer is “*History, dear boy, history*” to paraphrase a **disputed quote** of dear Harold (Macmillan that is).

In fact, both are **Domains**, and both are **DNA binding**. The illogicality of them being recorded in different places is accepted, however, to fix this early mistake now would not, it is claimed, be trivial. So, we live with it. So doing, click on the appropriate **Add** button and then prepare to head for the checkout desk (Good Grief! I am beginning to get used to this!).

Shimmy back to the top of the page. You should have **Basket 3** things in your basket.

Entry	Entry name	Organism	Remove
<input type="checkbox"/> P26367	PAX6_HUMAN	Homo sapiens (Human)	
<input type="checkbox"/> P26367[4-130]	PAX6_HUMAN	Homo sapiens (Human)	
<input type="checkbox"/> P26367[210-269]	PAX6_HUMAN	Homo sapiens (Human)	

Align BLAST Map Ids Download Clear Full View

Click on the basket to view your booty.

For each of the 3 items in turn (not all at once or you get all sequences in one file), select and **Download**.

Download selected (1)
 Download all (3)

Format:
 FASTA (canonical)

Compressed Uncompressed

Each time ensure you have **Download selected set** and the download parameters are set to **Uncompressed and FASTA (canonical)**. Then click the **Go** button.

The next few steps, as before, are very browser/OS dependant. Just do whatever it takes to save the three sequences in files called, as appropriate:

pax6_human.fasta
pax_domain.fasta
homeobox_domain.fasta

Now move back to America to the NCBI view of the **PAX6** gene. If you have problems getting there ... click here.

genomic	S70307.1	AAB30692.2
genomic	U63833.1 (3496..3772)	None
genomic	Z83307.1 (105..19232)	None
genomic	Z95332.1 (12642..20874)	None
mRNA	AB209177.1	BAD92414.1

Items 1 - 25 of 65
 << First < Prev Page 1 of 3 Next > Last >>

Near the bottom of the page, there is a section called **Related sequences**. Move to the bottom of the first page (of three) of the list of sequences. Click on the entry for the mRNA called **AB209177.1**. You will be rewarded by a **GenBank** entry in **GenBank** format.

Formats are tedious, but we will discuss them briefly at some point. You have already seen **FASTA** format. We will bump into **EMBL** format at some point. The other 137 or so formats are to be ignored!

Can you see the official gene name **PAX6**, mentioned in this entry? The **Gene Name** field (where **PAX6** should most certainly be mentioned) is entirely missing! If you searched **GenBank** (or **EMBL** come to that) for this sequence using the most obvious search **Keyword**, that is **PAX6**, do you think you would find this **PAX6** mRNA? You clearly should! A case for more consistent annotation? Perhaps something to consider further when we superficially mention the **Gene Ontology** project later.

Next, search the **Nucleotide** databases, by textual **Keyword**, for **PAX6** related sequences and download one or two for investigation. To achieve this worthy goal, move to the top of the current page and ensure that the database selection is set to **Nucleotide**. Click on the **Advanced** search option button.

Then in the **Nucleotide Advanced Search Builder**, change **All Fields** to **Title** in the pull down menu associated with the first search field and type in the keywords:

chromosome 11

In the second search field, again change **All Fields** to **Title** and type in the keyword:

paired box 6

You are asking **Entrez** to search for all **Nucleotide** database entries that contain the terms “**chromosome 11**” and “**paired box 6**” in the section of their annotation intended to be a succinct brief description (I.e. **Title**) of the entry. Click on the **Search** button to start the search going.

There is just one matching entry which is arrayed before you in **Genbank** format, very neat!! It was the **DEFINITION** line that you searched by selecting the **Field** value **Title**. I needed a few tries to get the right search to find just what was needed, and was a bit surprised at the simplicity and accuracy of the final search. You are looking at a **RefSeqGene** (a subset of the **RefSeq** database) entry. As such, it represents a genomic sequence for a “well-characterised gene”, in this case **PAX6**.

```
LOCUS       NG_008679                40170 bp    DNA     linear   PRI 23-DEC-2018
DEFINITION Homo sapiens paired box 6 (PAX6), RefSeqGene (LRG_720) on
            chromosome 11.
```

Take a look at the **FEATURES** for this entry. You will see that there are **three** genes mentioned. **PAX6**, of course. Also, on the strand that is the complement of that represented here, there is **PAX6-AS1** and **ELP4**.

```
gene       complement(<1..6396)
            /gene="PAX6-AS1"
            /gene_synonym="DKFZp686K1684"
            /note="PAX6 antisense RNA 1"
            /db_xref="GeneID:440034"
            /db_xref="HGNC:HGNC:53448"
gene       5001..38170
            /gene="PAX6"
            /gene_synonym="AN; AN2; ASG05; D11S812E; FVH1; MGDA; WAGR"
            /note="paired box 6"
            /db_xref="GeneID:5080"
            /db_xref="HGNC:HGNC:8620"
            /db_xref="MIM:607108"
```

Can you find the additional genes **PAX6-AS1** and **ELP4** in the genome displays you have looked at so far?

```
gene       complement(38437..>40170)
            /gene="ELP4"
            /gene_synonym="AN; AN2; C11orf19; dJ68P15A.1; hELP4;
            PAX6NEB; PAXNEB"
            /note="elongator acetyltransferase complex subunit 4"
            /db_xref="GeneID:26610"
            /db_xref="HGNC:HGNC:1171"
            /db_xref="MIM:606985"
```

```
join(16551..16560,20128..20258,21186..21401,22106..22271,
28174..28332,28848..28930,29160..29310,29409..29524,
32102..32252,32943..33028)
/gene="PAX6"
/gene_synonym="AN; AN2; D11S812E; FVH1; MGDA; WAGR"
/note="isoform a is encoded by transcript variant 1;
paired box protein Pax-6; paired box homeotic gene-6;
oculorhombin; aniridia type II protein"
/codon_start=1
/product="paired box protein Pax-6 isoform a"
/protein_id=" NP_000271.1 "
/db_xref="CCDS: CCDS31451.1 "
/db_xref="LRG:p1"
/db_xref="GeneID: 5080 "
/db_xref="HGNC: HGNC:8620 "
/db_xref="MIM: 607108 "
/translation="MQNSHSGVNLGGVFNVRPLPDSTRQKIVELAHSGARPCDISR
ILQVSNCGVSKILGRYYETGSIRPRAIGGSKPRVATPEVVSFKIAQYKRECPISFAWEI
RDRLLSEGVCCTNDNIPSVSSINRVLRLNLAASEKQMGADGMYDKLRMLNGQTGSWGTRP
GWYPGTSVPGQPTQDGCQQQEGGGENTNSISSNGEDSDEAQMRLQLKRLQRNRTSFT
QEQIEALEKEFERTHYPDVFAERLAAKIDLPEARIQWFNSRRRAKWRREEKLRNQR
QASNTPSHIPISSSFSTSVYQPIPOPTTPVSSFTSGSMLGRDOTALNTYSALPPMPS
FTMANNLPMQPPVPSQTSYSYSCMLPTSPSVNGRSYDTYTPPHMQTHMNSQPMGTSGETT
STGLISPGVSVVQVPGSEPDMSQYWPRLQ"
```

At the top of your page, Analyse **this sequence** by clicking on the **Highlight Sequence Features** option. The **CoDing Sequence (CDS)** feature for **PAX6** is displayed for you by highlighting the relevant parts (the coding **exons**) of the sequence and displaying the **CDS** details including the DNA regions that need to be **joined** to form the **CDS** and the **translation** of the **CDS**.

Use the controls at the bottom of your page to look at the other features of this entry (select feature **number** and then click on the **Feature** button).

What were the features that you found?

Why might you have expected more features than there were?

COMMENT REVIEWED [REFSEQ](#): This record has been curated by NCBI staff in collaboration with Isabel Hanson, David FitzPatrick. The reference sequence was derived from [Z95332.1](#) and [Z83307.1](#). This sequence is a reference standard in the [RefSeqGene](#) project.

PRIMARY	REFSEQ_SPAN	PRIMARY_IDENTIFIER	PRIMARY_SPAN	COMP
	1-18852	Z95332.1	2023-20874	
	18853-40170	Z83307.1	105-21422	

Take a look at the **COMMENT** and **PRIMARY** sections just above the **FEATURES**. This entry is suggested to be constructed from the alignment of two sequences from **GenBank**. The two aligned sequences being “**contigs**”, that is products of two individual sequencing projects of separate portions of the **PAX6** genomic region. We should discuss role of “**contigs**” in the human genome project, a little.

Take a quick look at the **GenBank** entries by entering their **ACCESSION** numbers (be sure to include the “.1”, the version number, at the end to avoid unwanted hits) into the **Search** box at the top of your page. Click on the **Search** button.

Nucleotide **Advanced**

Lo and behold, the two **GenBank** entries are summoned forth. Take a look at one or both. Not particularly illuminating I think². These are clones sequenced as part of the **Human Genome Project (HGP)**. They served to cover regions of **Chromosome 11** and have little biological significance in themselves.

- ✓ [Human DNA sequence from clone CFAT5 on chromosome 11, complete sequence](#)
- 1. 20,874 bp linear DNA
Accession: Z95332.1 GI: 2190397
[Taxonomy](#)
[GenBank](#) [FASTA](#) [Graphics](#)
- ✓ [Human DNA sequence from clone A1280 on chromosome 11, complete sequence](#)
- 2. 22,253 bp linear DNA
Accession: Z83307.1 GI: 1730464
[Taxonomy](#)
[GenBank](#) [FASTA](#) [Graphics](#)

Move back to the list, as illustrated. Select both entries.

Elect to **Analyse these sequences**, selecting from the extensive range of possibilities **Run BLAST**.

We will look at **blast** properly later, the idea here is to simple prove that these two sequencing clones really do overlap in the fashion suggested by the evidence so far. So, elect to **Align two or more sequences**³.

Cut and paste one of the sequencing clone **accession numbers** from the **Enter Query Sequence** box to the **Enter Subject Sequence** section of the form. Elect to **Show results in a new window**⁴. Firmly address the **BLAST** button.

Just one region of overlap should be identified.

Query	20771	GATCCGGAGCGACTTCCGCCTATTTCCAGAAATTAAGCTCAAACCTTGACGTGCAGCTAGT	20830
Sbjct	1	GATCCGGAGCGACTTCCGCCTATTTCCAGAAATTAAGCTCAAACCTTGACGTGCAGCTAGT	60
Query	20831	TTTTATTTTAAAGACAAATGTCCAGAGGGCTCATCATATTTTCCC	20874
Sbjct	61	TTTTATTTTAAAGACAAATGTCCAGAGGGCTCATCATATTTTCCC	104

How does the alignment you generated match up with the annotation of the original **RefSeq** entry you discovered?

- The annotation is very sparse which makes these entries very hard to find directly. The **EML-Bank** versions include some links to **Ensembl** codes. These would have been helpful but are not part of the official **International Nucleotide Sequence Database Collaboration (INSDC)** annotation that should be consistent between **GenBank**, **European Nucleotide Archive (ENA)**, which includes **EML-Bank**, and **DNA Data Bank of Japan (DDBJ)**.
- As opposed to comparing each of the two clones against an entire sequence database.
- Just because its neater. In my, significantly less than humble, opinion anyway.

Now for an entirely new search. The easiest way to get a fresh start is to move back to your browser tab displaying the GenBank Search results, and then click on the **Advanced** option of the **Search** facility at the top of the page. You should arrive back at the **Nucleotide Advanced Search Builder** offering a fresh start.

Set up a new search as illustrated and set it going. Ultimately simple this time. You have requested all **Human** sequences that are centrally associated with the gene **PAX6**.

A list of **60** or so sequences, all clearly claiming **PAX6** association and announcing their humanity loudly in Latin, will tumble forth.

You will have more hits than are displayed in one go, by default. Also, the hits are arranged in a “**Default**” order which has thus far defied all my attempts to associate with any reasonable definition of logic!

To deal with both of these issues, use the display control pull down menus at the top of your page to set the **items per page** to something big and the **Sort by** option to something sane.

The list shows matches between the terms entered and the **annotation** of DNA sequences. Not all relevant sequences will be present. For example, the **mRNA** with accession number **AB209177** was justifiably referenced in the **PAX6 Gene** entry but will not be in this list. **PAX6** appears nowhere in the annotation of **AB209177** including its **DESCRIPTION** (or **Title**) field.

Move far down the list, you will come to the **RefSeq PAX6 mRNAs** of a few pages back. Just before these entries is **M77844.1**. Save this one for later analysis. I choose **M77844.1** as it includes a few variations that will add interest. **Select the target sequence.**

You could now use the diminutive **Send to:** button which is near the bottom of your page to download all the selected sequences into a single file.

However, as there is only one sequence, and it would be so nice to be introduced properly before such intimacies as “downloading”. **Click on the link to the database entry** to see it in all its **GenBank Format** glory.

The sequence is for analysis rather than decoration, so use the **format menu** at the top of the page (currently set **GenBank**), and ask for **FASTA** format.

Now click the tiny **Send to:** button and **Choose Destination** to be **File**.

Strike the **Create File** button with a firm resolve. With irritating presumption, the choice of file name is made for you. Your sequence will be stored in a file named:

sequence.fasta

The **NCBI** is justifiably not famed for its understanding of poetry! **Do whatever it takes** to rename this file to be called:

pax6_mrna.fasta

One last file to save. Move back to your list of hits and deselect the mRNA that you have already saved.

Near the top of the list you should find two primer sequences. Their **Descriptions** suggest they are a pair of **PCR** primers used for picking out the **PAX6** gene. **Select both** by clicking in their selection boxes.

- [Homo sapiens neuroretina-specific pax6 gene enhancer region](#)
- 7. 267 bp linear DNA
Accession: AJ009907.1 GI: 3378599
[GenBank](#) [FASTA](#) [Graphics](#)
- [Homo sapiens paired box gene 6 \(PAX6\), isoform a sense primer](#)
- 8. 25 bp linear DNA
Accession: AJ270357.1 GI: 9557932
[GenBank](#) [FASTA](#) [Graphics](#)
- [Homo sapiens paired box gene 6 \(PAX6\), isoform a antisense primer](#)
- 9. 26 bp linear DNA
Accession: AJ270358.1 GI: 9557933
[GenBank](#) [FASTA](#) [Graphics](#)
- [Homo sapiens paired box protein PAX6 \(PAX6\) mRNA, complete cds](#)
- 10. 1,399 bp linear mRNA
Accession: AY047583.1 GI: 15422112
[GenBank](#) [FASTA](#) [Graphics](#)

```

LOCUS       AJ270357                25 bp    DNA     linear   PRI 26-JUL-2000
DEFINITION Homo sapiens paired box gene 6 (PAX6), isoform a sense primer.
ACCESSION   AJ270357
VERSION     AJ270357.1   GI:9557932
KEYWORDS    .
SOURCE      Homo sapiens (human)
  ORGANISM  Homo sapiens
            Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
            Mammalia; Eutheria; Euarchontoglires; Primates; Haplorrhini;
            Catarrhini; Hominidae; Homo.
REFERENCE   1 (bases 1 to 25)
  AUTHORS   Palm,K., Salin-Nordstrom,T., Levesque,M.F. and Neuman,T.
  TITLE     Fetal and adult human CNS stem cells have similar molecular
            characteristics and developmental potential
  JOURNAL   Brain Res. Mol. Brain Res. 78 (1-2), 192-195 (2000)
  PUBMED   10891600
REFERENCE   2 (bases 1 to 25)
  AUTHORS   Palm,K.
  TITLE     Direct Submission
  JOURNAL   Submitted (04-OCT-1999) Surgery, Cedars Sinai Medical Center, 8700
            Beverly Blvd., Los Angeles, CA 90048, US
COMMENT     Related entry: NM_000280.
FEATURES             Location/Qualifiers
     source           1..25
                     /organism="Homo sapiens"
                     /mol_type="genomic DNA"
                     /db_xref="taxon:9606"
     misc_feature     1..25
                     /note="PCR sense primer for paired box gene 6 (PAX6),
                     isoform a"
ORIGIN          1 ccagccagag ccagcatgca gaaca
//
    
```

Click on the **sense primer**. Properly, you would read all the **References** carefully. Instead, note the length looks about right and return to your list with the **Back** button.

It will be good to investigate these primers later, so find the diminutive **Send to:** button which is at the top of your page and use it. **Choose your Destination** to be **File** and set the **Format** of that file to be **FASTA**. Strike the **Create File** button with a confident click of your every ready mouse. Once more, the choice of file name is made for you. Your sequences are stored in a file named:

sequence.fasta

Do whatever it takes to rename this file to be called:

pax6_primers.fasta

- Complete Record
- Coding Sequences
- Gene Features

Choose Destination

- File Clipboard
- Collections

Download 2 items.

Format

FASTA

Sort by

Accession

Show GI

Create File

Back to **Ensembl**. More with the objective of looking at more sources of information via **Ensembl** than becoming expert **Ensembl** users.

Go to the **Ensembl** home page (www.ensembl.org). Choose to **View full list of all Ensembl species** using the link just under the **Select a species** menu.

Common name	Scientific name	Taxon ID	Ensembl Assembly	Accession	Genebuild Method	Variation database	Regulation database	Pre assembly
Aardvark (Pre)	<i>Orycteropus afer</i>	1230840	-	-	-	-	-	OrvAfe1@
Agassiz's desert tortoise	<i>Gopherus agassizii</i>	38772	ASM289641v1	GCA_002896415.1	Full genebuild	-	-	-
Algerian mouse	<i>Mus spretus</i>	10096	SPRET_EU_v1	GCA_001624865.1	External annotation import	-	Y	-
Alpaca	<i>Vicugna pacos</i>	30538	vicPac1	-	Projection build	-	-	-
Amazon molly	<i>Poecilia formosa</i>	48698	Poecilia_formosa-5.1.2	GCA_000485575.1	Full genebuild	-	-	-

Note that **Ensembl** (and **MapMaker**, of course) offers far more than just the **Human Genome**.

In particular, note the links to **EnsemblPlants**, **EnsemblFungi**, **EnsemblBacteria** etc. **Ensembl** databases at the bottom of the list.

Zebrafish	<i>Danio rerio</i>	7955	GRCz11	GCA_000002035.4	Full genebuild	Y	Y	-
Zig-zag eel	<i>Mastacembelus armatus</i>	205130	fMasArm1.1	GCA_900324485.1	Full genebuild	-	-	-

[Credits page for species images](#)

Other Metazoa

Additional metazoan genomes (initially insect vectors and nematodes) are available from [EnsemblMetazoa@](#)

Plant and fungal genomes can be found at [EnsemblPlants@](#) and [EnsemblFungi@](#)

Protists, Bacteria and Archaea

Unicellular eukaryotic and prokaryotic genomes can be found at [EnsemblProtists@](#) and [EnsemblBacteria@](#) respectively.

During this exercise, you will only look at the **Human genome**, by far the most completely recorded. However, all the other **Ensembl** genomes are behind the same interface. The techniques required to examine the Human genome are broadly those required to examine any **Ensembl** genome.

Move back to the **Ensembl** home page and go to the **Human PAX6** gene information by setting the **Search** fields as shown and clicking the **Go** button boldly.

Search

Human ▼ for

PAX6 **Go**

e.g. [BRCA2](#) or [rat 5:62797383-63627669](#) or [rs699](#) or [coronary heart disease](#)

The target gene is at the top of the hit list.

Click on the link to the **PAX6 (Human Gene)**.

You should recognise the view you now see. The list of transcripts and view of the genomic region exactly as you examined via the **NCBI**.

PAX6 (Human Gene)
ENSG00000007372 11:31784779-31818062;-1
 Paired box 6 [Source:HGNC Symbol;Acc:HGNC:8620]
 LRG_720 (LRG display in Ensembl gene record; description: Locus Reference Genomic record for **PAX6**,) is an external reference matched to Gene ENSG00000007372
[Variant table](#) • [Phenotypes](#) • [Location](#) • [External Refs.](#) • [Regulation](#) • [Orthologues](#) • [Gene tree](#)

PAX6-202 (Human Transcript)
ENST00000379107 11:31789194-31810305;-1
 Paired box 6 [Source:HGNC Symbol;Acc:HGNC:8620]
 P26367 (UniProtKB/Swiss-Prot record; description: Paired box protein **Pax-6**) is an external reference matched to Translation ENSP00000368401
[Location](#) • [External Refs.](#) • [cDNA seq.](#) • [Exons](#) • [Variant table](#) • [Protein seq.](#) • [Population](#) • [Protein summary](#)

PAX6-203 (Human Transcript)
ENST00000379109 11:31788911-31810667;-1
 Paired box 6 [Source:HGNC Symbol;Acc:HGNC:8620]
 P26367 (UniProtKB/Swiss-Prot record; description: Paired box protein **Pax-6**) is an external reference matched to Translation ENSP00000368403
[Location](#) • [External Refs.](#) • [cDNA seq.](#) • [Exons](#) • [Variant table](#) • [Protein seq.](#) • [Population](#) • [Protein summary](#)

PAX6-204 (Human Transcript)
ENST00000379111 11:31789946-31811952;-1
 Paired box 6 [Source:HGNC Symbol;Acc:HGNC:8620]
 AOA1W2PQA8 (UniProtKB/TrEMBL record; description: Paired box protein **Pax-6**) is an external reference matched to Translation ENSP00000368406
[Location](#) • [External Refs.](#) • [cDNA seq.](#) • [Exons](#) • [Variant table](#) • [Protein seq.](#) • [Population](#) • [Protein summary](#)

There is much to investigate here, but maybe that should wait for a specialised **Ensembl** course. They are run regularly in **Cambridge** and elsewhere.

To make a bit more space, elect to **Hide transcript table**.

This gene has 84 transcripts ([splice variants](#)), [145 orthologues](#), [50 paralogues](#), is a member of [2 Ensembl protein families](#) and is associated with [30 phenotypes](#).

Transcripts, Orthologues and Paralogues.

At the top of the page, note the summary giving, particularly, an expectation of the numbers of

Begin by taking a look at how **Ensembl** sees the **Homologues** of **PAX6**. First the **Orthologues** and then the **Paralogues**. Click on the **Orthologues** link in the left hand side of your browser page.

Take a look at some of the alignments providing support for the homologous relations. The protein alignments are the more informative (from the **View Sequence Alignments** menu, select **View Protein Alignment**).

Armadillo (<i>Dasyus novemcinctus</i>)	1-to-1 View Gene Tree	PAX6 (ENSDNOG0000000761) Compare Regions (JH561443) View Sequence Alignments	Orthologue Alignment View Protein Alignment View cDNA Alignment
---	--	--	---

Which human **PAX6** isoform has been chosen to align with **orthologues**?

How do you suppose this choice might have been justified?

At the bottom of your screen, **Ensembl** offers a list of organisms with no **PAX6 Orthologue**.

Species without orthologues
9 species are not shown in the table above because they don't have any orthologue with ENSG00000007372
<ul style="list-style-type: none"> Monterrey platyfish (<i>Xiphophorus couchianus</i>) <i>Ciona intestinalis</i> Dog (<i>Canis lupus familiaris</i>) <i>Caenorhabditis elegans</i> (<i>Caenorhabditis elegans</i>) Lamprey (<i>Petromyzon marinus</i>) <i>Saccharomyces cerevisiae</i> (<i>Saccharomyces cerevisiae</i>) <i>Ciona savignyi</i> Fruitfly (<i>Drosophila melanogaster</i>) Alpaca (<i>Vicugna pacos</i>)

Can you comment of the inclusion of **Drosophila** in this list?

Once your curiosity concerning **Orthologue** alignments is completely sated, click on the **Paralogues** link.

All **50** that **Ensembl** expected (see above) are listed in a seemingly randomised series. This is not very helpful.

What regions of **PAX6** would you expect might have **Paralogues** (or **Orthologues**, come to that)?

In order to easily make sense of this list, rank it by some measure of **Quality**, click on the **Query %id** column header as many times as it takes to achieve an ordering of the list of **paralogues** that is **High** → **Low** by **Query %id**.

You should now be able to discern at least **2** distinct sets of **paralogues** by looking down the **Ensembl identifier and gene name** column.

At the top of the list you should find genes **paralogous** to the **Paired Box** domain of **PAX6**.

Type	Ancestral taxonomy	Ensembl identifier & gene name	Compare	Location	Target %id	Query %id
Ancient paralogues	Bilateral animals (Bilateria)	ENSG00000009709 PAX7 paired box 7 [Source:HGNC Symbol;Acc:HGNC:8621]	<ul style="list-style-type: none"> Region Comparison Alignment (protein) Alignment (cDNA) 	1:18,631,006-18,748,866:1	31.73 %	37.84 %
Ancient paralogues	Bilateral animals (Bilateria)	ENSG00000135903 PAX3 paired box 3 [Source:HGNC Symbol;Acc:HGNC:8617]	<ul style="list-style-type: none"> Region Comparison Alignment (protein) Alignment (cDNA) 	2:222,199,888-222,298,996:-1	31.68 %	36.70 %
Ancient paralogues	Bilateral animals (Bilateria)	ENSG00000106331 PAX4 paired box 4 [Source:HGNC Symbol;Acc:HGNC:8618]	<ul style="list-style-type: none"> Region Comparison Alignment (protein) Alignment (cDNA) 	7:127,610,292-127,618,114:-1	41.40 %	32.57 %
Ancient paralogues	Bilateral animals (Bilateria)	ENSG00000125618 PAX8 paired box 8 [Source:HGNC Symbol;Acc:HGNC:8622]	<ul style="list-style-type: none"> Region Comparison Alignment (protein) Alignment (cDNA) 	2:113,215,997-113,278,950:-1	30.44 %	31.42 %
Ancient paralogues	Bilateral animals (Bilateria)	ENSG00000174279 EVX2 even-skipped homeobox 2 [Source:HGNC Symbol;Acc:HGNC:3507]	<ul style="list-style-type: none"> Region Comparison Alignment (protein) Alignment (cDNA) 	2:176,077,472-176,083,913:-1	14.71 %	16.05 %
Ancient paralogues	Bilateral animals (Bilateria)	ENSG00000119614 VSX2 visual system homeobox 2 [Source:HGNC Symbol;Acc:HGNC:1975]	<ul style="list-style-type: none"> Region Comparison Alignment (protein) Alignment (cDNA) 	14:74,239,472-74,262,738:1	18.56 %	15.37 %
Ancient paralogues	Bilateral animals (Bilateria)	ENSG00000106410 NOBOX NOBOX oogenesis homeobox [Source:HGNC Symbol;Acc:HGNC:22448]	<ul style="list-style-type: none"> Region Comparison Alignment (protein) Alignment (cDNA) 	7:144,397,240-144,410,227:-1	9.70 %	15.37 %
Ancient paralogues	Bilateral animals (Bilateria)	ENSG00000052850 ALX4 ALX homeobox 4 [Source:HGNC Symbol;Acc:HGNC:450]	<ul style="list-style-type: none"> Region Comparison Alignment (protein) Alignment (cDNA) 	11:44,260,440-44,310,166:-1	16.06 %	15.14 %

Further down, list entries declare a **paralogous** association with the **Homeobox** domain of **PAX6**.

How many of the **PAX6 paralogues** are associated with the conservation of the **Paired Box** domain?

View some of the protein alignments between the gene **PAX6** and its **paralogues**.

Some **paralogues** seem to have two regions of high similarity (e.g. **PAX4** or **PAX2**), others only one (e.g. **PAX1**)? Can you explain?

Click on the **Domains & features** link (from **Transcript-based displays** → **Protein Information**).

Domain source	Start	End	Description	Accession	InterPro
PANTHER	34	434	FAMILY NOT NAMED	PTHR24329	-
PANTHER	34	434	PAIRED BOX PROTEIN PAX-6	PTHR24329:SF294	-
Gene3D	208	285	-	1.10.10.60	-
Prosite_profiles	222	282	HOMEODOMEX_2	PS50071	IPR001356 [Display all genes with this domain]
Smart	224	286	HOX_1	SM00389	IPR001356 [Display all genes with this domain]
Pfam	226	281	Homeobox	PF00046	IPR001356 [Display all genes with this domain]
CDD	226	283	homeodomain	cd00086	IPR001356 [Display all genes with this domain]
Prosite_patterns	257	280	HOMEODOMEX_1	PS00027	IPR017970 [Display all genes with this domain]
SuperFamily	6	143	Homeodomain-like	SSF46689	IPR009057 [Display all genes with this domain]
SuperFamily	205	283	Homeodomain-like	SSF46689	IPR009057 [Display all genes with this domain]
Pfam	4	142	PAX	PF00292	IPR001523 [Display all genes with this domain]
Smart	4	142	pax3	SM00351	IPR001523 [Display all genes with this domain]
Prosite_profiles	4	144	PAIRED_2	PS51057	IPR001523 [Display all genes with this domain]
CDD	5	145	PAX	cd00131	IPR001523 [Display all genes with this domain]
PRINTS	8	23	PAIREDBOX	PR00027	IPR001523 [Display all genes with this domain]
PRINTS	26	44	PAIREDBOX	PR00027	IPR001523 [Display all genes with this domain]
PRINTS	60	77	PAIREDBOX	PR00027	IPR001523 [Display all genes with this domain]
PRINTS	78	95	PAIREDBOX	PR00027	IPR001523 [Display all genes with this domain]
Gene3D	1	86	Winged helix-like DNA-binding domain superfamily	1.10.10.10	IPR036388 [Display all genes with this domain]
Gene3D	87	162	Winged helix-like DNA-binding domain superfamily	1.10.10.10	IPR036388 [Display all genes with this domain]

Are you surprised that the precise location of the **PAX6** Homeobox domain is not identically predicted by the **SMART** and **Pfam Domain Databases**? If not, why not?

How is that all the predictions, of different domain databases, for a **Paired domain** have the same **Interpro** identifier?

Why does **PRINTS** appear to predict four **Paired_domains**?

Click on the link to the **SMART** entry for the **Paired domain** (**SM00351**).

Here you will find (quoted from **Interpro**) a **Description** of a **Paired domain**.

Where would you expect a **Paired domain** to occur in a protein?


What expectations do you have concerning what typically follows a **Paired domain**?

The paired domain is an approximately 126 amino acid DNA-binding domain, which is found in eukaryotic transcription regulatory proteins involved in embryogenesis. The domain was originally described as the 'paired box' in the Drosophila protein paired (prd) [(PUBMED:2877747), (PUBMED:3123319)]. The paired domain is generally located in the N-terminal part. An octapeptide [(PUBMED:10811620)] and/or a homeodomain can occur C-terminal to the paired domain, as well as a Pro-Ser-Thr-rich C terminus.

Paired domain proteins can function as transcription repressors or activators. The paired domain contains three subdomains, which show functional differences in DNA-binding. The crystal structures of prd and Pax proteins show that the DNA-bound paired domain is bipartite, consisting of an N-terminal subdomain (PAI or NTD) and a C-terminal subdomain (RED or CTD), connected by a linker. PAI and RED each form a three-helical fold, with the most C-terminal helices comprising a helix-turn-helix (HTH) motif that binds the DNA major groove. In addition, the PAI subdomain encompasses an N-terminal beta-turn and beta-hairpin, also named 'wing', participating in DNA-binding. The linker can bind into the DNA minor groove. Different Pax proteins and their alternatively spliced isoforms use different (sub)domains for DNA-binding to mediate the specificity of sequence recognition [(PUBMED:11103953), (PUBMED:15148315)].


The second paragraph of the **Description** claims, in gross summary:

- A paired domain is a DNA binding domain that has 2 binding regions each of which involves a helical triplet
- The second and third helices of each helical triplet form **Helix-Turn-Helix (HTH)** motifs



- The **HTH** regions bind the **DNA major groove**⁵

- The first helical triplet is preceded by a **β-turn** and **β-hairpin** (“wing”) that participate in the DNA binding





- The linker region between the two helical triplets can bind the **DNA minor groove**

Bear this in mind when looking at the 3D structures a couple of pages on.

Click on **Display all genes with this domain** for the **Paired domain** and **Homeobox domain** **InterPro** families. The locations of all genes including each domain will be displayed graphically and textually. **PAX6** is shown in red.

Paired domain - IPR001523

Homeobox domain - IPR001356



Which domain, **Paired domain** or **Homeobox domain** is more common in humans?

How many human **PAX** genes are there?

Are all the **PAX** genes on **Chromosome 11**?

⁵ If, like me, you have conceptual problems with major and minor grooves. Try this [animated picture](#). Helped me at least. As did the image above.

Move back to the **Domains & features** display. Link to the **InterPro** database entry for **Paired domain**, also known as **IPR001523**. Here you will find the origins of the **SMART** documentation. Click on the **Proteins matched** link. You will see listed a number of representations of proteins that, according to **InterPro**, include a **Paired domain**. Amongst these will be the human **PAX6** protein, also known as **P26367**⁶.


P26367  	Paired box protein Pax-6	Homo sapiens (Human)	
---	--------------------------	----------------------	--

Click on the **Structures** link in the top left hand corner of the page. **InterPro** will offer links to relevant entries in the **PDBe**, **SCOP** and **CATH**⁷ databases. Click on the link to the **6pax** entry in the **PDBe** database. You will arrive at the entry for **6pax** in **PDBe**, the European version of **PDB** maintained at the **EBI**. Views of this structure are offered on the right hand side of the page. Click on the largest image which shows the paired box protein domain binding DNA rather beautifully. Once you have admired this image, in all its various guises, sufficiently, move back to the **6pax PDBe** entry. From the **Quick links** on the right of the page, select the **3D Visualisation** option.



The **SMART** documentation you read earlier suggested two paired box sub-domains, each of which “... form a three-helical fold, with the most C-terminal helices comprising a **helix-turn-helix (HTH)** motif that binds the **DNA major groove**”. Move your image around to confirm this assertion.

The same **SMART** documentation claims the sub-domain nearer the **N terminal** “... encompasses an N-terminal **beta-turn** and **beta-hairpin**, also named '**wing**', participating in **DNA-binding**. The linker can bind into the **DNA minor groove**”. Manipulate your image to investigate the veracity of these assertions.



Once you have seen all there is to see of **6PAX**, move back to the **Ensembl Domains & features** display. Try the same tricks with the **InterPro Homeobox domain**. This time, it is difficult to find **P26367** in the huge list⁸ **Proteins matched**, but you do not need to in order to link to the **Structures**. There are many more structures to choose from this time. I suggest you go for **2cue**. You have to imagine the DNA this time.

It looks rather as if the **Homeobox domain** also includes a helical triplet including a **Helix-Turn-Helix**. You could have confirmed this by reference to the relevant **SMART** documentation (as you did for the **Paired box domain**). It is the **HTH** that the **Homeobox** uses to bind to DNA.

InterPro did not detect the Homeobox HTH as it did the Paired box HTH. Have you any thoughts as to why this might be?

Can you explain the strangely frayed ends displayed in some of the representations of the **2cue** 3D structure?

6 Third from the bottom of the first page, last time I counted.

7 **PDB** is the main database for **3D** protein structures. **SCOP** and **CATH** are also **3D** structure related databases.

8 If you really wanted to, the best approach is to search for **P26367** in the search box at the top of the page and then look for the **Homeobox domain** entry in the **Detailed signature matches** list.

To end, a gesture towards demonstrating that you could quite easily have computed most of the information you have been accessing, ready packed, from various databases. There are many way this objective could be achieved, I choose to search for the features of the **PAX6** protein.

As has been discovered from several information sources, the **PAX6** human protein has two DNA binding domains. A paired box at the **N terminal** and a homeobox a little further along. Both of the domains include **Helix-Turn-Helix (HTH)** motifs. In this exercise, you will investigate how you might discover these domains and motifs using the various freely available domain databases (discussed previously) and other feature prediction programs. Clearly, this is superfluous for this particularly, well documented protein, but a valuable option in other circumstances.

One approach would be to consider each relevant domain database in turn. Each major domain database has its own Home web site and customised software to take **Query** protein sequences, compare those sequences with domain representations (typically based on **Hidden Markov Models**) and to report convincing matches. This would work, but would be tedious as there are many viable databases to consider. It would be dangerous to rely on too few of the databases available as none is perfect. You need a consensus prediction to be sure you miss nothing.

Also, you would need to know which databases are particularly appropriate for each domain you considered might be present. All databases cannot be optimised for all types of domain (for example, the **SMART** database specialises in domains that occur in signalling proteins).

So, let us not search individual domain databases. I am sure you could find your own way through using most of the major searches, if you wished. Notes on using the **Prosite**, **Pfam** and **PRINTS** domain databases appear in the discussion sections of appropriate exercises, but should not take up significant class practical time I feel. Investigating each individually turn does have some merit however. **Prosite** illustrates how widely domain matches can vary in significance, **Pfam** gives and opportunity to superficially discuss **HMMs** and searching **PRINTS** illustrates the small margin between a positive and a negative result.

Here, use just **Interpro** to do the whole job. **Interpro** will search for all domains using the appropriate domain databases, thus removing the tedium of interrogating a miscellany of domain searching resources individually.



Interpro defines protein families according to the way that proteins match elements of a wide range of protein family databases, including all those we have discussed thus far. **Interpro** provides a search tool that will search all or any of the major protein family databases and assign **Interpro** family associations to the query protein(s) accordingly. To have a look at some of the possibilities offered by **Interpro**, **Go to:**

<http://www.ebi.ac.uk/interpro/>

If you were to enter the **PAX6** human protein into the obvious place on the **InterPro** home page and click the **Submit** button, you would produce exactly the results you saw many pages back, when you were investigating **UniProtKB**⁹. **Do this** if you have the time and inclination.

By implication, **InterPro** offers a fuller experience via the **InterProScan** search tool. Other than the opportunity not to search **ALL** the domain databases, and having the results arranged slightly differently, I am unsure what the extra effort brings? Never mind, there are many things of which I am unsure, so, from the **InterPro** Home page ...

Tools | InterProScan



InterProScan is a sequence analysis application (nucleotide and protein sequences) that combines different protein signature recognition methods into one resource. [More about InterProScan](#)

Select the **InterProScan** link. Here you will be offered the opportunity to **download the InterProScan** program.

I am not sure this is too useful an offer for most? But it is there.

For now, **chose the online Sequence search**.

Sequence search


InterProScan sequence search
Click here to scan your protein sequence and discover the domains it contains and the family to which it belongs.

⁹ Not surprising as **UniProtKB** simply links to **Interpro** to show you its graphic.

You will arrive at a page very similar to that from which you started, as far as the offer to run a domain search is concerned? Except! We now have **Advanced options**. Click on the **Advanced options**.

The **Advanced options** only allow you to choose which databases you wish to search and which feature prediction programs you wish to run. The default is to use all the databases and to run all the predictor programs. I struggle to imagine an occasion I would want to save the **EBI** servers a few cycles by considering which options to deselect, but it so nice to know I could if I wished to.

In passing, the offer to run the feature predictor programs in the **Other sequence features** section is relatively new. Of course, all these programs could be run individually from their home websites (follow the links behind the program names), in the same way as the domain databases can be searched individually. **Interpro** just aims to make thing easy for the user. The programs currently offered are:




- **Coils** is a program for predicting **coiled coils**.
- **MobiDB Lite** is a method of **Fast and highly specific consensus prediction of intrinsic disorder in proteins**. A new facility for **Interpro**. It uses **MobiDB**, a database of annotations of **intrinsic protein disorder**. **Protein disorder** being a structural features characterising large sets of proteins with prominent members that are **intrinsically disordered proteins**.
- **Phobius** & **TMHMM** are programs to predict **Transmembrane regions** (essentially **hydrophobic, uncharged** regions). There is no reason to expect any **Transmembrane regions** in this protein.
- **SignalP** predicts the presence and location of **signal peptide cleavage sites** in amino acid sequences from different organisms. I am pretty certain that there is no reason to expect signal peptides in this protein.

Do you think it a good idea for **Interpro** to offer feature prediction programs as well as domain database searches?

Paste the human **PAX6** sequence into the patiently waiting box (from the file you made earlier called **pax6_human.fasta**). Accept the **“do everything”** default. Click on the **Submit** button.

After several moments of deep thought, filtering and validating, you will be presented with a table of results looking very much like the one your saw earlier when looking around **UniProtKB**.



Do you think the **Coil** prediction might be correct?

Notice that **Interpro** assigns both the **PAX** domain and the **Homeobox** domain of human **PAX6** to the **Interpro** family **Homeobox domain-like**. Both of these associations are based on the hit behind the link **SSF46689**.

SCOP classification	
Root:	SCOP hierarchy in SUPERFAMILY [SCOP_0] (11)
Class:	All alpha proteins [SCOP_46456] (284)
Fold:	DNA/RNA-binding 3-helical bundle [SCOP_46688] (14)
Superfamily:	Homeodomain-like [SCOP_46689] (19)
Families:	Homeodomain [SCOP_46690] (40)
	Recombinase DNA-binding domain [SCOP_46728] (5)
	Myb/SANT domain [SCOP_46739] (15)
	SLIDE domain [SCOP_100998]
	GARP response regulators [SCOP_81683]
	DNA-binding domain of telomeric protein [SCOP_46745] (2)
	Paired domain [SCOP_46748] (3)

Follow this link and you will see it leads to the **Homeodomain-like superfamily** of the  database that specialises in very general (**SCOP¹⁰ superfamily level**) protein classifications. One **Superfamily** entry will typically correspond to a number of more specific **SCOP** classifications. Here you can see that the **Superfamily** domain **Homeodomain-like** includes both the **Homeodomain & Paired domain Families**.

Return to your **Interpro** results page. The links beginning “**GD3SA**” point to **Superfamily** domains defined by the **CATH Protein Structure Classification database**. **CATH** is similar to **SCOP** in that it is another Structural classification database. **CATH Superfamilies** are to be found in the **Gene3D** database¹¹. One such link suggests two regions that belong to a **Winged helix-like DNA-binding domain superfamily**. These seem to correspond to the two **Helix Triplets** of the **Paired domain**. Note that the **Helix Triplet** in the **Homeobox domain** is not detected by **Gene3D**? Possibly because of the lack of **Beta Sheet “Wings”** in the **Homeobox domain**?

Interpro provides a unified report of all the superfamilies detected either by reference to the **SCOP** or **CATH** databases.



Click on the region bars and you will be offered links to the relevant **Interpro** entries.

Contributing signatures
Signatures from InterPro member databases are used to construct an entry.
GENE3D ⓘ
G3DSA:1.10.10.10 (G3DSA:1.10.10.10)

Follow one of the links to the **Interpro** family **Winged helix-like DNA-binding domain superfamily (IPR036388)**. Note the **Contributing signatures** in the top right hand corner of the page. Here is listed the domain database entries that are used to determine the presence of an **Interpro Winged helix-like DNA-binding domain superfamily**

Essentially, if **GENE3D** finds a match with its **Winged helix-like DNA-binding domain superfamily (G3DSA:1.10.10.10)**, then **Interpro** records a match with its **Winged helix-like DNA-binding domain superfamily (IPR036388)**.

Contributing signatures
Signatures from InterPro member databases are used to construct an entry.
SUPERFAMILY ⓘ
SSF46689 (SSF46689)

Move back to your **Interpro** graphic and follow one of the links to the **Interpro** family **Homeodomain-like domain superfamily (IPR009057)**. Again, note the **Contributing signatures**.

This time it is stated that, if **Superfamily** finds a match with its **Homeodomain-like superfamily (SSF46689)**, then **Interpro** records a match with its **Homeodomain-like domain (IPR009057)**¹².

I conclude the **Homologous superfamilies** and **Domains and Repeats** sections of the graphic simply summarise and confirm information from the **Detailed signature matches** section.

¹⁰ Structural Classification Of Proteins.

¹¹ Broadly, **CATH** is to **Gene3D** as **SCOP** is to **Superfamily**.

¹² Until recently, matches with **Gene3D** entries were also regarded as significant here.

While you have the **Interpro Homeobox-like domain superfamily** in view, it is easy to obtain an impression of how widely spread throughout nature is this domain family. You have already established that there are a fair few in human proteins.

Click on the **Species** button on the left hand side of the page.

As you can see, this is a very popular domain. By clicking on the appropriate  button, you can get to either the protein sequences in **Fasta** format or list their accessions codes. Try a few, but be careful! It really does get you **ALL** the sequences, and that is often quite a lot, which can take time.

Homologous Superfamily
Species: Homeobox-like domain superfamily
 (IPR009057)
Key Species

Key species	Number of proteins	FASTA	Protein IDs
<i>Arabidopsis thaliana</i> (Mouse-ear cress)	1277		
<i>Homo sapiens</i> (Human)	1074		
<i>Danio rerio</i> (Zebrafish)	919		
<i>Oryza sativa</i> subsp. japonica (Rice)	908		
<i>Mus musculus</i> (Mouse)	860		
<i>Drosophila melanogaster</i> (Fruit fly)	464		
<i>Caenorhabditis elegans</i>	225		
<i>Escherichia coli</i> (strain K12)	94		
<i>Saccharomyces cerevisiae</i> (strain ATCC 204508 / S288c) (Baker's yeast)	31		
<i>Schizosaccharomyces pombe</i> (strain 972 / ATCC 24843) (Fission yeast)	24		

Taxa

- cellular organisms 964386 proteins | FASTA | Protein IDs
- Archaea 2945 proteins | FASTA | Protein IDs
- Bacteria (eubacteria) 792252 proteins | FASTA | Protein IDs
- Eukaryota (eucaryotes) 169189 proteins | FASTA | Protein IDs
- unclassified sequences 3998 proteins | FASTA | Protein IDs
- Viruses 636 proteins | FASTA | Protein IDs
- other sequences 10 proteins | FASTA | Protein IDs

Proteins matched: Homeobox-like domain superfamily (IPR009057)
 'Filtered by species: *Schizosaccharomyces pombe* (strain 972 / ATCC 24843) (Fission yeast) (excludes child species) (change species)

Showing 1 to 20 of 24 results

Accession	Protein name	Species	Domain architecture
O13719	SWIRM domain-containing protein lat1	<i>Schizosaccharomyces pombe</i> (strain 972 / ATCC 24843) (Fission yeast)	
O13788	SWI/SNF and RSC complexes subunit ssr1	<i>Schizosaccharomyces pombe</i> (strain 972 / ATCC 24843) (Fission yeast)	
O14013	RNA polymerase I-specific transcription initiation factor rrm5	<i>Schizosaccharomyces pombe</i> (strain 972 / ATCC 24843) (Fission yeast)	
O14108	DNA-binding protein eta2	<i>Schizosaccharomyces pombe</i> (strain 972 / ATCC 24843) (Fission yeast)	

You can make this list enormous by injudicious employment of the expansion buttons (the **Number of protein** links). Why not? It amused me for a few moments anyway.

Finally, return again to you **Interpro** graphic. Notice that the **Paired domain** prediction is supported by matches with **six** different domain databases. Only **four** of these support the **Homeobox domain** prediction. The missing two database matches are with **Prosite patterns** (identifier begins **PS** and typically matches the domain partially where it is best conserved) and with **PRINTS** (identifier begins **PR**).

IPR001523 Paired domain

- PS00034 (PAIRED_1)
- PS51057 (PAIRED_2)
- SM00351 (ps3)
- cd00131 (PAX)
- PF00292 (PAX)
- PR00027 (PAIREDBOX)

IPR001356 Homeobox domain

- SM00389 (HOX_1)
- PS50071 (HOMEBOX_2)
- cd00086 (Homeobox)
- PF00046 (Homeobox)

IPR017970 Homeobox, conserved site

- PS00027 (HOMEBOX_1)

Why do you suppose there is no match from **PRINTS** or **Prosite patterns** to support the **Homeobox domain** prediction for this protein?

What do you suppose the **Homeobox conserved site** might be?

THE END

DPJ – 2019.01.30

Model Answers to Questions in the Instructions Text.

Notes:

For the most part, these “**Model Answers**” just provide the reactions/solutions I hoped you would work out for yourselves. However, sometime I have tried to offer a bit more background and material for thought? Occasionally, I have rambled off into some rather self indulgent investigations that even I would not want to try and justify as pertinent to the objective of these exercises. I like to keep these meanders, as they help and entertain me, but I wish to warn you to only take regard of them if you are feeling particularly strong and have time to burn. Certainly not a good idea to indulge here during a time constrained course event!

Where things have got extreme, I am going to make two versions of the answer. One starting:

Summary:

Which has the answer with only a reasonably digestible volume of deep thought. Read this one.

The other will start:

Full Answer:

Beware of entering here! I do not hold back. Nothing complicated, but it will be long and full of pedantry.


This makes the Model answers section very big. **BUT**, it is not intended for printing or for reading serially, so I submit, being long and wordy does not matter. Feel free to disagree.

From your investigations using **UniProtKB**:

Describe the arrangement of Helices within **PAX6**.

From the evidence of the textual table and the graphic, there are **nine** helices in all, that occur in groups of **three**.

Aligning the graphical representation of the positions of these helices with the **Interpro** domain prediction graphics (discovered via **UniProtKB** earlier), it is clear that the first two of the helical triplets lie in the **Paired** domain and the third is in the **Homeobox** domain



From your investigations using **Entrez**:

What were the features that you found?

Summary:

The first feature was the **CoDing Sequence (CDS)** for a **PAX6** isoform, the canonical isoform a. The **NCBI** say they omit the other isoform(s) as they do not aspire to “completeness” but just an indication of structure with the **RefSeq** entries.

The other three features were the coding sequences for three **ELP4** isoforms. Why more than one for this gene then? Possibly because they are “more different” representing interesting variation in gene structure?

```
complement(39424..>39569)
/gene="ELP4"
/gene_synonym="AN; AN2; C11orf19; dJ68P15A.1; hELP4; PAX6NEB; PAXNEB"
/inference="similar to AA sequence (same species):RefSeq:NP_001275654.1"
/exception="annotated by transcript or proteomic data"
/note="isoform 2 is encoded by transcript variant 2; elongator complex protein 4; PAX6 neighbor gene protein; elongation protein 4 homolog"
/codon_start=3
/product="elongator complex protein 4 isoform 2"
/protein_id=" NP_001275654.1 "
/db_xref="CCDS: CCDS73271.1 "
/db_xref="GeneID: 26610 "
/db_xref="HGNC: HGNC:1171 "
/db_xref="MIM: 606985 "
/translation="MAAVATCGSVAASTGSAVATASKSNVTSFORRGRPRASVTNDSGPRLVSIAGTRPSVRNGQLLVSTGLPALDQLLGGGLAVGTVLLIEEDKYNIVSPLLKFKYLAEGIVNGHTLLVASAKEDPANILQELPAPLLDDCKCKEFEDVYVNHKTPESNIKMKIAMRYQLLPKMEIGPVSSSRFGHYDASKRMPQELIEASNWHGFFLPEKISSTLKVPECSLTPGYTKLLQFIQNIIEEFGDGSNPKQKQRNLRIGIQNLGSPWGGDDICCAENGGNSHSLTKFLYVLRGLRRTLSLACIITMPTHLIQNKAIARVTTLSDVVVGLESFVIGSERETNPLYKDYHGLIHRQIPRNLNLCDESDVKDLAFKLRKLFTEIAGVQNHDLGSRRTNIPYPPGFSYLLKQKDSAWGEGSLQHSFTLMSFLAKATAFASRLVRHSEPLKQNGSGRIRQAAGPRLWHBGRQRREAPGLLGIPP"
```

```
complement(39438..>39569)
/gene="ELP4"
/gene_synonym="AN; AN2; C11orf19; dJ68P15A.1; hELP4; PAX6NEB; PAXNEB"
/inference="similar to AA sequence (same species):RefSeq:NP_061913.3"
/exception="annotated by transcript or proteomic data"
/note="isoform 1 is encoded by transcript variant 1; elongator complex protein 4; PAX6 neighbor gene protein; elongation protein 4 homolog"
/codon_start=1
/product="elongator complex protein 4 isoform 1"
/protein_id=" NP_061913.3 "
/db_xref="CCDS: CCDS7875.2 "
/db_xref="GeneID: 26610 "
/db_xref="HGNC: HGNC:1171 "
/db_xref="MIM: 606985 "
/translation="MAAVATCGSVAASTGSAVATASKSNVTSFORRGRPRASVTNDSGPRLVSIAGTRPSVRNGQLLVSTGLPALDQLLGGGLAVGTVLLIEEDKYNIVSPLLKFKYLAEGIVNGHTLLVASAKEDPANILQELPAPLLDDCKCKEFEDVYVNHKTPESNIKMKIAMRYQLLPKMEIGPVSSSRFGHYDASKRMPQELIEASNWHGFFLPEKISSTLKVPECSLTPGYTKLLQFIQNIIEEFGDGSNPKQKQRNLRIGIQNLGSPWGGDDICCAENGGNSHSLTKFLYVLRGLRRTLSLACIITMPTHLIQNKAIARVTTLSDVVVGLESFVIGSERETNPLYKDYHGLIHRQIPRNLNLCDESDVKDLAFKLRKLFTEIARHLHLPDLSDTVSRSSKMDLAESAKRLGPGCGMAGGKXHLDF"
```

```
complement(39533..>39569)
/gene="ELP4"
/gene_synonym="AN; AN2; C11orf19; dJ68P15A.1; hELP4; PAX6NEB; PAXNEB"
/inference="similar to AA sequence (same species):RefSeq:NP_001275655.1"
/exception="annotated by transcript or proteomic data"
/note="isoform 3 is encoded by transcript variant 3; elongator complex protein 4; PAX6 neighbor gene protein; elongation protein 4 homolog"
/codon_start=2
/product="elongator complex protein 4 isoform 3"
/protein_id=" NP_001275655.1 "
/db_xref="CCDS: CCDS73272.1 "
/db_xref="GeneID: 26610 "
/db_xref="HGNC: HGNC:1171 "
/db_xref="MIM: 606985 "
/translation="MAAVATCGSVAASTGSAVATASKSNVTSFORRGRPRASVTNDSGPRLVSIAGTRPSVRNGQLLVSTGLPALDQLLGGGLAVGTVLLIEEDKYNIVSPLLKFKYLAEGIVNGHTLLVASAKEDPANILQELPAPLLDDCKCKEFEDVYVNHKTPESNIKMKIAMRYQLLPKMEIGPVSSSRFGHYDASKRMPQELIEASNWHGFFLPEKISSTLKVPECSLTPGYTKLLQFIQNIIEEFGDGSNPKQKQRNLRIGIQNLGSPWGGDDICCAENGGNSHSLTKFLYVLRGLRRTLSLACIITMPTHLIQNKAIARVTTLSDVVVGLESFVIGSERETNPLYKDYHGLIHRQIPRNLNLCDESDVKDLAFKLRKLFTEIAGVQNHDLGSRPQRLGSSNSPASASLVAGITGAHHHTQLIFVFLVEMGFHHVQAGLELLTSGDSSASQASGITGMSYRARPRLYFKENKSKVGAROLLETREEHLSRLLILTQAEELCMGRFFFTAFHIFNELPCKGDCICLCTQCTQ"
```

Full Answer:

Note that only the final coding exon of **ELP4** is within this **RefSeq** sequence, which is defined as the genomic region for **PAX6**. This is clear from the length of the **translations** offered. The exon referenced is only long enough to code for just over **40** amino acids which is far shorter than any of the three entire isoform sequences offered here.

Note also that this final coding exon of **ELP4** (stretching from **39424/39438/39533** to **39569** of this **RefSeq** entry) does **not** overlap the coding region of the **PAX6** gene itself (stretching from **16551** to **33028** of this **RefSeq** entry).

In fact, the two genes do not overlap according to the evidence. The **PAX6** gene extends from **5001** to **38170**. The portion of the **ELP4** gene that is included in this entry extends from **40170** (the end) to **38437** (in the opposite direction). This give a gap between the two genes stretching from **38171** to **38436**.

RefSeqGenes, comprise the entire gene plus **5,000** “extra” base pairs in either direction. The overlap here is entirely within the “extra” base pairs.

```
join(16551..16560,20128..20258,21186..21401,22106..22271,28174..28332,28848..28930,29160..29310,29409..29524,32102..32252,32943..33028)
/gene="PAX6"
/gene_synonym="AN; AN2; D11S812E; FVH1; MGDA; WAGR"
/note="isoform a is encoded by transcript variant 1; paired box protein Pax-6; paired box homeotic gene-6; oculorhombin; aniridia type II protein"
/codon_start=1
/product="paired box protein Pax-6 isoform a"
/protein_id=" NP_000271.1 "
/db_xref="CCDS: CCDS31451.1 "
/db_xref="LRG:p1"
/db_xref="GeneID: 5080 "
/db_xref="HGNC: HGNC:8620 "
/db_xref="MIM: 607108 "
/translation="MQNSHSGVNLGGVFNVRPLPDSTRQKIVELAHSGARPCDISRILQVSNCGVSKILGRYYETGSRIPRAIGGSKPRVATPEVSKIAQYKRECPISFAWEIRDRLLESEGCTNDNIPVSSINRVLRLNLAASEKQMGADGMYDKLRMLNGQTSWGRTPGWYPTGSPVGPQPTQDGCQQEQEGGENTNKSISNGEDSDEAQMRLQLKRLQRNRTSFTQEIQEIALEKEFERHYDPVFAERELAAKIDLPEARIQVWFNSRRAKWRREEKLRNQRRQASNTPSHIPISSSFSVSYQPIPOPTTPVSSFTSGSMLGRDTRDALTNTYSALPPMPSFTMANNLPMQPPVPSQTSYSSYCLPTSPVSNGRSYDTYTPPHMQTHMNSQPMGTSGETSTGLISPGVSVVQVPGSEPDMSQYWRPLQ"
```

```
gene 5001..38170
/gene="PAX6"
/gene_synonym="AN; AN2; D11S812E; FVH1; MGDA; WAGR"
/note="paired box 6"
/db_xref="GeneID: 5080 "
/db_xref="HGNC: HGNC:8620 "
/db_xref="MIM: 607108 "
```

Careful study of either of the two **Genome Database** displays visited earlier (**Genome Data Viewer** or **Ensembl**) will confirm the relative positions of **PAX6** and **ELP4**.

```
gene complement(38437..>40170)
/gene="ELP4"
/gene_synonym="AN; AN2; C11orf19; dJ68P15A.1; hELP4; PAX6NEB; PAXNEB"
/note="elongator acetyltransferase complex subunit 4"
/db_xref="GeneID: 26610 "
/db_xref="HGNC: HGNC:1171 "
/db_xref="MIM: 606985 "
```

The annotation (specifically the **gene_synonyms**) of **ELP4** associate this gene with **PAX6**. However, as the **ELP4** gene annotation to the right attests, only because of its proximity.

General protein information

Preferred Names
elongator complex protein 4

Names
PAX6 neighbor gene protein
elongation protein 4 homolog

Why might you have expected more features than there were?

Summary:

All the evidence has suggested that **PAX6** has at least **2** isoforms. This would lead me to expect at least **2** CDS features here related to **PAX6**?

Full Answer:

The explanation from the **NCBI** is that this sort of **RefSeq** entry is intended to be used as a template against which sequences from an individual can be mapped to seek variations. Only a token **CDS** feature is included to indicate the position of the gene. For such an entry, recording every isoform is not essential.

This sounded convincing to me, Until I began to wonder why there were three **CDS** features for **ELP4** which is not even the gene primarily represented by this entry? Maybe I will ask more questions if and when I ever have the strength. In the meantime, mostly for my information, I record their exact explanation here.

“ ... note that **RefSeqGene** defines genomic sequences to be used as reference standards for well-characterized genes. These sequences serve as a stable foundation for reporting mutations, for numbering exons and introns, and for defining the coordinates of other variations. We normally select one **RefSeq** transcript to serve as a reference standard. The goal is not to record all introns and exons of all isoforms, but just to choose one representative to help define the locus. Therefore, most of our **RSG** records have only a single **RefSeq** as reference standard. If an **LSDB** manager or other stakeholder requests that other **RefSeqs** be added as alternate standards, this can easily be done (with the complication that, if a public **LRG** exists, the **RefSeqGene** record is fixed). We receive requests from stakeholders to include **RefSeqs** that represent all known exons, or **RefSeqs** that have become community standards. Often, after creating an **RSG** using our own internal criteria, we receive stakeholder requests to change or add transcripts. Many of these requests come from the **LRG** project regarding transcripts to be included on the **LRG** records.

Generally, **RefSeq** accessions can be added or removed without reversioning, unless a transcript is upgraded or a new one defined that extends beyond the bounds of the **RSG**, or matches a new build of the genome, in which case the **RSG** will be extended and reversioned as needed.

Regarding the chromosomal locus, our standard range is 5 kb upstream from the 5' end and 2 kb downstream from the 3' end of the mRNAs with the greatest extent. For this calculation, we do indeed use all available **RefSeq** (NM_) accessions. If the database manager or stakeholder has information on promoters or other upstream or downstream regulatory regions, we can certainly extend the **RefSeqGene** locus to accommodate these.

Regarding mismatches, the goal is to exactly match the current build of the genome, unless there is overwhelming transcript and EST evidence that a mismatch should be retained.

Regarding the confusing subject of exon numbering, exon numbers are currently provided only on **RSG** genomic records based on a subset of available transcript **RefSeqs** for the gene. These are often those selected by locus-specific databases as reference sequence reporting standards. You can find an explanation of how exons are numbered here:

<http://www.ncbi.nlm.nih.gov/refseq/rsg/faq/#exon>

You will find links to more information on **RefSeqGenes** on the home page for the **RefSeqGene** project:


<http://www.ncbi.nlm.nih.gov/refseq/rsg/>

Regarding the **PAX6 RSG** sequence, only difference I see between **NG_008679.1** and the current build of the genome (**GRCh38**) is an extra 'G' beyond the 3'-UTR of the **PAX6** transcripts (at **NC_000011.10:g.31,819,125**). ... “

Yes, well I think I followed most of that? and that my interpretation is broadly correct? In summary, there are no fixed rules.

How does the alignment you generated match up with the annotation of the original RefSeq entry you discovered?

The most intuitive way of encapsulating graphically the way these two sequencing clones overlap was donated by **Cecilia Pinto (Oeiras, 2013.12.09-12)**. Thank you Cecilia.



From your investigations using **Ensembl**:

Which human **PAX6** isoform has been chosen to align with **orthologues**?

How do you suppose this choice might have been justified?

Species	Gene ID	Peptide ID	Peptide length	% identity (Protein)	% coverage	Genomic location
Human (<i>Homo sapiens</i>)	ENSG00000007372	ENSP00000492024	436 aa	92 %	94 %	11:31784779-31818062
Anole lizard (<i>Anolis carolinensis</i>)	ENSACAG00000002252	ENSACAP00000002317	422 aa	95 %	97 %	1:60705219-60743090


```

CLUSTAL W (1.81) multiple sequence alignment

ENSP00000492024/1-436      MQNSHSGVNQLGGVFNVRPLPDSTRQKIVELAHSGARPCDISRILQTHADAKVQLDNQ
ENSACAP00000002317/1-422 MQNSHSGVNQLGGVFNVRPLPDSTRQKIVELAHSGARPCDISRILQ-----
                        *****

ENSP00000492024/1-436      NVSNGCVSKILGRYYETGSIRPRAIGGSKPRVATPEVVS KIAQYKRECPSIFAW EIRDRL
ENSACAP00000002317/1-422 -VSN GCVSKILGRYYETGSIRPRAIGGSKPRVATPEVVS KIAQYKRECPSIFAW EIRDRL
                        *****

```

The protein used to represent **PAX6** human is consistently **ENSP00000492024**. At least, this was the choice for the alignments I looked at. This is the protein sequence of **isoform 5a** (as evidenced by the extra **14** amino acids around the end of the first line), probably chosen as it is the longer option (**436** amino acids as opposed to **422**) and so (from the crude informatics viewpoint) represents more information.

The canonical isoform is used for the non-human sequence, I imagine because that is what you get if you pull the sequence from **UniProt** (or similar). If I am right, more computational convenience than Biological justification.

Can you comment of the inclusion of **Drosophila** in this list?

The claim that there are no **Human PAX6 Orthologues** for **Drosophila** does seem a bit strange even entirely wrong?) given the abundance of evidence to the contrary available from the **INTERNET** and beyond. For example:

Source 1, Source 2, Source 3, Sources 4, Source 5 ...

However, the people I speak to from **Ensembl** are adamant, so maybe I am missing something. I suspect a more accurate statement might be:

*“There is insufficient evidence for the **Ensembl** automatic **Orthology** detection procedures to identify any **Drosophila Orthologues** for **Human PAX6**.”*

What regions of **PAX6** would you expect might have **Paralogues** (or **Orthologues**, come to that)?

It has been established that **PAX6** has **2** domains. A **Paired Box** domain and a **Homeobox** domain.

It is surely reasonable to expect that **Homologues** of **PAX6** will correspond to either one of these domains? Or possibly to both?

The list of **50** is not organised in any way that makes it easy to decide the domain(s) to which each **Paralogue** might be associated, Some manipulation is required for a clear view.


In this exercise, ranking by quality (defined by alignment **Sequence Identity**) is used. This works because the **Paired Box** matches are much longer than the **Homeobox** matches. I would not expect this approach to work in all cases however. Maybe there should be a way to separate the types of **Paralogues** in a more reliable fashion?

What are the first two bases and what are the last two bases of nearly every intron?

As you are probably well aware, introns are highly conserved at each end. They typically begin with **GT** and end with **AG**. This rule is obeyed by all but one of the introns of this transcript (**intron 3-4** starts **GC** rather than **GT**).

As the cartoon suggests, the conservation does not apply just to the first and last two bases, but that is where the conservation is most strict. So strict that when exceptions from this rule were sought in the databases, it was thought most of the deviations were due to annotation error!

The cartoon also suggests that introns have **C/T rich regions** towards their ends (the **Polypyrimidine tract**). This too is clearly evident in most of the introns of this transcript, even though only small parts of the introns are displayed.



How long is the sixth exon and why would this concur with your expectations?

It is **42** base pairs long, so it codes for **14** amino acids. Specifically, it codes for the **14** extra amino acids that define **isoform 5a**.

Explain the **Start Phase** and **End Phase** columns?

An exon/intron boundary can occur anywhere in a codon. The **Start** and **End Phases** record how an intron has been inserted into a coding region with respect to the coding reading frame.

If an exon ends at the end of a codon, then its **End Phase** is **0**.

Clearly, the next exon must begin at the start of a codon. Its **Start Phase** is also **0**.


If an exon ends after the first base of a codon, then its **End Phase** is **1**.

Clearly, the next exon must begin after the first base of a codon. Its **End Phase** is also **1**.

If an exon ends after the second base of a codon, then its **End Phase** is **2**.

Clearly, the next exon must begin after the second base of a codon. Its **End Phase** is also **2**.

I attempt a picture, though I am sure that is clear? I just like pictures, and lots of colours.



Why does **Prints** appear to predict four **Paired_domains**?

Prints does not find the **Homeobox_domain** at all. If you were to investigate by using the **Prints** search carefully, you will find it nearly does, but the evidence is not quite strong enough. As has been discussed, none of these systems are perfect. They all occasionally fail. That is why it is always best to use **Interpro** to consult them all and deliver a consensus answer.

Prints appears to find **FOUR Paired_domains**. This is only because of the way **Prints** works. **Prints** finds **FOUR** signatures (or **motifs**) that together indicate **ONE Paired domain**. **Prints** searches for ordered series of **motifs** that together indicate **domains**. Here it reports each of four motifs separately, but it is only claiming one **Paired domain**.

PRINTS	8	23	Paired domain	PR00027	IPR001523 [Display all genes with this domain]
PRINTS	26	44	Paired domain	PR00027	IPR001523 [Display all genes with this domain]
PRINTS	60	77	Paired domain	PR00027	IPR001523 [Display all genes with this domain]
PRINTS	78	95	Paired domain	PR00027	IPR001523 [Display all genes with this domain]

Which domain, **Paired domain** or **Homeobox domain** is more common in humans? How many human **PAX** genes are there?

As you will have expected, there are but **9 Paired domains** in the Human genome. There are many more **Homeobox domains**. Note particularly that **Interpro** predicts far more **Homeobox** domains than **Ensembl** admits to. **Ensembl** predictions are based purely on computer searches and comparisons, which can never be entirely perfect.

Are all the **PAX** genes on **Chromosome 11**?

Of course not? What a stupid question!

Well, I suppose they could all be on **Chromosome 11**? By chance ... or maybe design ... who knows, the lack of predictable pattern in all this business never ceases to astound me.


But, philosophy aside, the answer is **NO**.

Can you explain the strangely frayed ends displayed in some of the representations of the **2cue** 3D structure?

2cue is a 3D structure determined by Nuclear Magnetic Resonance (NMR). This is a process that does not involve immobilizing the target as a crystal (as is the case with structures determined by **X-ray crystallography**). Parts of the protein will still be moving around whilst its structure is being determined.

I think of **NMR** as analogous to taking a long exposure photograph of a group of children. Each child will appear in many different places! The frayed ends represent various positions in which the ends of the **homeobox** were detected during the **NMR** process.

In some views, including the one you were offered to move around, all the possible positions are averaged out before the structure is stored. I prefer the fuzzy view ... much more fun.



I broadly believe that which I have just typed, however, I must stress that my understanding of **NMR** is tragically incomplete. If anyone would like to offer a better explanation, I am very willing to hear it.

From your investigations of Domain & Motif identification using Interpro

Do you think it a good idea for **Interpro** to offer feature prediction programs as well as domain database searches?

Well ... why not? The purpose of **InterProScan** is to associate regions of query proteins with **Interpro** domains. This was originally achieved, exclusively, by simply comparing a query sequence with all entries of relevant individual domain databases. These entries being representations of alignments of examples of specific domains constructed by homology searching (i.e. **blast** and similar).

I would suggest including a few predictor programs would provide extra evidence gathered from more general, more theoretical definitions of domains. I would imagine the inclusion of these programs has improved and widened the picture provided by **InterProScan**.

Searching domain databases, typically composed of **HMM profiles**, such as **Pfam**, **Prosite** and **PRINTS** is quite different to running the predictor programs. As I cannot improve on the justification of this claim offered to me by Geoff Barton (Head of the group responsible for **Jalview**, **Jpred**, **Jnet** and much more), I will just reproduce his explanation here:

“ ... The main difference is that with an **HMM profile** you have a "specific" example of a domain or motif whereas with something like **COILS**, you have something trained across all examples.

For example, for secondary structure prediction, you could (a) do predictions of alpha-helix and beta-strand just by aligning a sequence to a protein of known structure, or an **HMM** from a family of aligned proteins of known structure. This is a specific case of secondary structure in the context of one protein family. Or (b) you can train a predictor from **ALL** protein families and then apply this. The advantage of (a) is it is very specific to the individual protein family and so should be more accurate for that family. The disadvantage is that it does not generalise to proteins that are not very like the specific example. The advantage of (b) is that it will work with any protein but will likely be less accurate than (a) for proteins that fit into the (a) category. ... “

Do you think the Coil prediction might be correct?

I do not recall anything in what we have discovered thus far that would directly suggest there should be a **coiled coil** here, in the middle of the **HTH**. However, **wikipedia** does suggest **coiled coils** are associated with **transcription factors** (which **pax6_human** is).

" ... Many **coiled coil**-type proteins are involved in important biological functions such as the regulation of **gene expression**, e.g. **transcription factors**. ... "

I think I would not be overly convinced by this prediction, but I would not make that judgement with any great confidence. The all knowing **wikipedia** says:

“ ... **Coiled coils** usually contain a repeated pattern, **hxxhxc**, of **hydrophobic (h)** and **charged (c)** amino-acid residues, referred to as a **heptad repeat**. ... “

Geoff Barton comments:

“ ... Sometimes the pattern that is particular to **coiled-coils** also turns up in other helices that pack against each other. You would need to look at some examples of coiled-coil structures to see if the example you are using fits structurally. ... “

Which seems very reasonable. The **heptad repeat** pattern could easily occur just by chance. **COILS** surely cannot predict the structure of the helices well enough to make an assured judgement? **COILS** offers a suggestion the user must follow up with other resources.

There is also the evidence that **Jpred** (a system for secondary structure prediction that you will meet later), possibly using the **COILS** program disguised as **LUPAS**, does not detect any coiled coils. This could be for a number of reasons. Possibly **LUPAS** is not the same program as **COILS**, or it is a different version, or **Jpred** runs **COILS**, but with different parameters.

Not many clear and confident answers in Bioinformatics are there!

Discussion Points and Casual Questions arising from the Instructions Text.**Notes:*****Work in progress I fear.***

The intention is to provide a full consideration of some issues skimmed over in the exercise proper.

If you are attending a “supervised” presentation of the exercise, I would hope to have conducted a live discussion of all these issues to an extent that reflects:

- the depth that seems appropriate
- the time available
- the degree to which the issues seem to match the interests of the class
- how many of you are awake

Here, I hope to write out very full answers were such a response exists. Accordingly, I suggest you will not need to read much of many of these discussions. There will be much detail of interest to rather few of you. Possibly a bit self indulgent, but I wish to make a note of all the background I have discovered while writing these exercises.

In a nutshell, the exercises are trying to make very general points avoiding too much detail. Nevertheless, I record the detail outside the main exercise text, just in case it might be of interest. Some of the answers to the “**Casual Questions**” are exceedingly trivial. Some of the “**Discussion Points**” are exceedingly long and rambling. You have been warned.

Can you now say how many transcripts there are according to the **Genome Data Viewer**?

11, count the transcript prediction lines of blobs and wiggly lines.

Discussion of the **Ensembl** transcript colour and numbering schemes.

Introduce Ensembl pipeline

Introduce Vega ... for a number of vertebrates

Havana = group feeding Vega for human/Mouse and similar ... not all genomes of Vega

GENCODE ... amalgamation of Vega and ensembl pipeline ... source of Ensembl transcript predictions

Conclusion: gold ... agreed between pipeline and Vega

red ... either Vega or pipeline, used to be able to tell which by the transcript number (≥ 200 pipeline, < 200 Vega) but now all numbers 200+

blue ... non-protein coding

The naming/numbering of transcripts is being improved. Current temporary. Future a method representing prediction quality.

Source ... Latest gems from Ben of Ensembl (Email 2017.09.25)

Strategies employed to minimise the time spent on searches employed to determine gene structures, specifically with respect to their implementation by **Ensembl**.

As described already, assuming a suitable comprehensive set of appropriate sequences, the location and structure of all transcripts could be determined by a simple two stage operation:

mapping all quality mRNA sequence onto the genome to discover the

In particular:

first genscan ... find most genes

then CCDS (CDS agreed by pipeline, Vega and NCBI ... Human/Mouse specific at present) search on genscan hits only reveals coding regions accurately

then mRNA (RefSeq and other high quality data/predictions) only on CCDS hits ... reveals UTRs accurately

Why it is reasonable to not regard a match of a **RefSeq** mRNA with the **Genome** as, by itself, sufficient evidence to uniquely predict a transcript.

RefSeq mRNA sequences are not determined by careful sequencing of individual mRNA/cDNA. If they were, it would be difficult to argue with the **NCBI** approach of regarding a quality match between a **RefSeq** mRNA and the genome as sufficient evidence to predict the location of a transcript.

However, **RefSeq** mRNA sequences are actually computed from assemblies of many single pass, poor quality, cDNA sequences (**ESTs**).

Ensembl regards these sequences as good evidence but not conclusive by themselves.

NCBI appears to rely more on the reliability of **RefSeq** mRNA sequences.

How reliable would you judge these predictions to be?

Needs thought and investigation here but ... main message is that there is huge variance in quality between these predictions! Far from binary announcement of existence or otherwise.

Specifically, there is only **APPRIS** support where there is **CCDS** matching. This makes sense as a **CCDS** hit implies a relationship to a confident protein isoform that is very likely to have **orthologues**. This will make more sense when we have considered how many **PAX6** isoforms there might be.

Sequence formats, specifically FASTA format.

Indeed, sequence formats will be discussed, but a little further down. Until then, try to contain thy urgent thirst for elucidation.


Discussion of the isoform alignments.

Not much to say? ... see the inserted 14 amino acids in the middle of the PAX domain?

Refer to silly domain / DNA Binding confusion, although I think I do that elsewhere.

Can you see the evidence for this assertion in the regional genomic maps of a few pages back?

Yep ... it is visible in the **Genome Data Viewer** version. The default **Ensembl** pictures are too crushed.




Are the **Interpro** results broadly as you might expect?

Yep ... two domains, **homeo** & **PAX**, as suggested by **NCBI**. Here more domain/motif databases are quoted, but the conclusion is the same both sides of the Atlantic.

Note the inconsistent naming of the domains! Is this really necessary one muses? Life is muddled enough already surely. How long does it take to choose a single name?

This graphic will be considered in more detail later when we look at **Interpro** more closely.



Sequence formats.

2 varieties required. For analysis (**FASTA**) or for storage in a database with annotation (**GenBank**, **EMBL**).

FASTA for all the sequences saved so far, minimal annotation, just enough for identification. The sequence is the issue.

```
>NAME Description
Sequence ... ..
>NAME Description
Sequence ... ..
```

Genbank or **EMBL** (why two!?) where the annotation is the primary focus (although a bit silly without the sequence!). Formats for the databases. Pity there is two, but to expect too much sanity between **EMBL** and **NCBI** is clearly asking too much. Here we look at **Genbank**, later we will see **EMBL**. I will not elaborate, both have online manuals (**GenBank**, **EMBL**). The basics are intuitive (I hope).

Some reference to the times of many many formats here???

Can you see the official gene name **PAX6**, mentioned in this entry?

No ... **PAX6** occurs several times in the page (try searching with **Ctrl F**) but only in the page annotation, not in the databases entry!

Do you think you would find this **PAX6** mRNA using the search term **PAX6**?


Absolutely not!!

A superficial mention of the **Gene Ontology Project**.

Very superficially ... possibly here?

Can you find the additional genes **PAX6-AS1** and **ELP4** in the genome displays you have looked at so far?

The **Genome Data Viewer** picture prefers to show the location of the **RefSeq** mRNAs that support the existence of **ELP4**.



Hover of any of these and the link to **ELP4** is revealed.

ELP4

Gene: ELP4
 Title: elongator acetyltransferase complex subunit 4
 Location: 31,509,729..31,784,525
 Length: 274,797
 Merged features: NP_001275655.1 and NM_001288726.1

Download: [NP_001275655.1](#), [NM_001288726.1](#)

The **Ensembl** display you viewed earlier clearly includes all its predictions of transcripts for **ELP4**.

The **Genome Data Viewer** picture shows the location of the **RefSeq** RNA that support the existence of **PAX6-AS1**.

DKFZp686K1684

Gene: DKFZp686K1684
 Title: uncharacterized LOC440034
 Location: 31,816,566..31,887,041
 Length: 70,476

ncRNA: NR_033971.1
 Title: uncharacterized LOC440034
 Location: 31,816,566..31,887,041
 [Length]
 Span: 70,476
 Placed: 1,656
 Product: 1,656

Download: [NR_033971.1](#)

Hover over the RNA reference. An association with a gene called “**DKFZp686K1684**” is revealed. But

gene complement(<1..6396)
 /gene="PAX6-AS1"
 /gene_synonym="DKFZp686K1684"
 /note="PAX6 antisense RNA 1"
 /db_xref="GeneID:440034"
 /db_xref="HGNC:HGNC:53448"

“**DKFZp686K1684**” is the gene-synonym of **PAX6-AS1**. So the gene is discovered, if indirectly.

This **gene synonym** implies that this gene was originally identified by the **German Cancer Research Centre (DKFZ)**.

No mention of **PAX6-AS1** Though? Unless you **Download NR_033971.1** and look at the **FASTA** description line which reads:

```
>gi|300068930|ref|NR_033971.1|:1-1656 Homo sapiens PAX6 antisense RNA 1 (PAX6-AS1), long non-coding RNA
```

Declaring that this **RefSeq RNA** is a feature of a non-coding gene called **PAX6-AS1**. The only reason for naming it such being that it slightly overlaps the **PAX6** gene on its antisense strand.

PAX6-AS1 is also represented in the **Ensembl** view of the **PAX6** region. However, it is not so easy to find. Certainly there is no obvious evidence is the view as you examined it previously.

To find **PAX6-AS1** (even disguised as **DKFZp686K1684**), should you really want to, try the following. First add the **RefSeq human import** track to your display. To achieve this, elect to **Configure this page**. In the **Genes** subsection of the **Genes and transcripts** section, turn on **RefSeq human import**, choosing **Expanded with labels**.

Enable/disable all Genes

Comprehensive Gene Annotations from GENCODE 27

Basic Gene Annotations from GENCODE 27

CCDS set

EST-based

RefSeq GFF3 annotation import


RefSeq human import

Finally click on the in the top right hand corner to **Save and close** your selections.

RefSeq human import.	NR_033971.1 > misc RNA
NM_001288726.1 > protein coding	<input type="checkbox"/> NR_misc
NM_001288725.1 > protein coding	<input type="checkbox"/> NR_misc
NM_019040.4 > protein coding	<input type="checkbox"/> NR_misc
XM_005252865.2 > protein coding	<input type="checkbox"/> NR_misc

Essentially, you have asked for some **RefSeq** based predictions from the **NCBI** to be added to the display. Amongst these (top right) is the misc RNA prediction based on the match between the **RefSeq** sequence **NR_033971.1**

Job done? Well ... yes I think so, but having travelled so far! Let us proceed to the tortuous end.



Notice the suspicious matching of some of the **RCN1** gene transcripts and the new **misc RNA** from the NCBI (I removed the **Genes (Basic set from GENCODE 27)** track from the display to enhance clarity).

For a closer look, adjust the Region in view to expose the whole of that covered by **NR_033971.1**


To do this

Go to Region In Detail for more tracks and navigation options (e.g. zooming)

and then blunder about with the

zooming and **sliding** and **dragging** tools until you get a view something like mine ... **OR**, you could use the trick it has just taken me an hour to discover, of selection the area desired by entering the **RCN1** gene name in the appropriate place!!!

Location: 11:31812391-32105755 Go
Gene: RCN1 Go



Now I am even more convinced by the similarity of structure of the **processed transcript RCN1-202** and the **misc RNA NR_033971.1**

Name	Transcript ID	bp	Protein	Biotype	CCDS	UniProt	RefSeq	Flags
RCN1-201	ENST0000054950.3	2572	331aa	Protein coding	CCDS7876	Q15293 V9HW95	NM_002901 NP_002892	TSL:1 GENCODE basic APPRIS P2
RCN1-210	ENST00000532942.5	1191	280aa	Protein coding	-	Q15293	-	TSL:2 GENCODE basic APPRIS ALT2
RCN1-204	ENST00000528630.1	549	28aa	Protein coding	-	H0YDA4	-	CDS 5' incomplete TSL:3
RCN1-206	ENST00000530348.5	527	58aa	Protein coding	-	E9PP27	-	CDS 3' incomplete TSL:4
RCN1-209	ENST00000532721.1	494	19aa	Protein coding	-	E9PLM2	-	CDS 3' incomplete TSL:3
RCN1-203	ENST00000527337.1	737	57aa	Nonsense mediated decay	-	H0YER5	-	CDS 5' incomplete TSL:3
RCN1-202	ENST00000506388.2	1658	No protein	Processed transcript	-	-	NR_033971	TSL:1
RCN1-208	ENST00000532474.5	714	No protein	Processed transcript	-	-	-	TSL:3
RCN1-205	ENST00000530146.1	635	No protein	Processed transcript	-	-	-	TSL:3
RCN1-207	ENST00000531345.1	2653	No protein	Retained intron	-	-	-	TSL:2
RCN1-211	ENST00000533898.5	2416	No protein	Retained intron	-	-	-	TSL:2

Time to look at the transcript table for **RCN1** for the detail.

Go to the **RCN1 Ensembl** gene page using the main search option at the top of your current page. Make sure the transcript table is in view and take a look at the entry for the **processed transcript RCN1-202**.

By the Lord Harry! **processed transcript RCN1-202** is based exclusively on the evidence of the match between **NR_033971.1** and the genome!

I conclude that what the NCBI predict as the **non-coding gene PAX6-AS1** with a single transcript based upon the **RefSeq** RNA **NR_033971.1**, **Ensembl** predicts as a non-coding transcript of the protein coding gene **RCN1**.


All this nonsense achieves rather little, in the context of the exercise, I suppose. I certainly do not want to suggest you follow your way through the pain I have just endured. However, I hope this little diversion into pedantry does illustrate how using multiple sources of imperfect information can be less than straight forward. To obtain a complete picture often requires lot of effort and patience. Never mind, it will ever get better ... possibly.

The role of “contigs” in the human genome project.

The objective here is to establish some understanding of what these two sequences that you have found are. To do this it is necessary to understand how the Human Genome was determined using the sequencing technologies available at the turn of the century.

Broadly, the **Human Genome** was considered too big to sequence in one step. Each **Chromosome** was therefore processed separately.

However, even the smallest **Human Chromosome** was too large to be efficiently sequenced as a single entity. Accordingly, **Chromosomes** were fragmented randomly into manageable sections (**20-40Mb** at the start of the project, up to **150Mb** by the end). Each fragment was cloned and sequenced separately. The sequences determined for the chromosome fragments are, in this context, referred to as **Contigs**. The **Contigs**, once reassembled, determined the sequence of each entire **Chromosome**. Time for President Clinton to, somewhat optimistically, announce the task completed.



All the individual **Contig** sequences are retained in specialist databases. A minimal selection of the **Contigs** are stored in more general databases such as those you are searching in this exercise. The selected **Contigs** form a “**Golden Path**” through the assembly of all **Contigs**. The “**Golden Path**” is such that the entire **Chromosome** is represented using the smallest set of **contigs** practical.


Clearly, just the **contigs** of the “**Golden Path**” would be insufficient to reliably determine the **Full Chromosome Sequence**. “**Golden Path**” elements might overlap only by a few tens of base pairs. Such an overlap would not be credible except for the knowledge that it is supported by many other **contigs** stored elsewhere.

So, you are looking at the two “**Golden Path**” **contigs** whose overlap fully encompasses the entire **PAX6** gene. Your next task is to use blast to compute the overlap between the two **contigs**.

To conclude, a final note on the term “**Contig**”.

CONTIG

An overlapping set of Sequencing Reads



Contig (short for **Contiguous**) was a term introduced by Rodger Staden to mean an overlapping set of sequencing reads.

Once assembled, any overlapping set of sequencing reads (**Contig**) will acquire a **Consensus Sequence** that is its single best representation.

The ultimate objective of any sequencing project is to create a single **Contig** that represents the entire target region. The **Consensus Sequence** of this final **Contig** will be “**The Answer**”.

Inevitably, due to incomplete data and/or insufficiently clever software, the initial assemblies generated many partial region **Contigs**. Sequencing and assembling must continue until a whole region **Contig**, of acceptable quality, is computed.

For reasons of convenience, the term **Contig** has come to mean the **Consensus sequence** associated with a **Contiguous** set of sequencing reads. This is the meaning I have used in the preceding discussion.

How many of the **PAX6** paralogues are associated with the conservation of the **Paired Box** domain?

The first **8** entries in the “*Quality Ordered*” list of **Paralogues** are recorded as being associated with the **Paired Box** domain of **PAX6** in the **Ensembl identifier and gene name** column. So there are **9 PAX** paralogues for Human (according to **Ensembl**, and all the other sources I have come across). They are **PAX1 PAX2 PAX3 PAX4 PAX5 PAX6 PAX7 PAX8** and, last but by no means least, **PAX9**.

The remaining **42** list entries are recorded as **paralogous** to the **Homeobox** domain of **PAX6**.

Are you surprised that the precise location of the **PAX6** Homeobox domain is not identically predicted by the **SMART** and **Pfam Domain Databases**? If not, why not?

Both **Smart** and **Pfam** predict the locations of

Smart	224	286	Homeobox domain	SM00389	IPR001356	[Display all genes with this domain]
Pfam	226	281	Homeobox domain	PF00046	IPR001356	[Display all genes with this domain]

protein domains. They both use similar, but not identical, methods. In this case, both predict a **Homeobox domain** where it is very likely that there is a **Homeobox domain**. This is surely very good news. Should we really expect the predicted locations to be identical? These are just predictions after all and it is questionable whether domains really have precise amino acid specific locations. It is doubtful that all human experts would agree on the most probable exact domain location. Why would we expect computer programs to do better?

How is that all the predictions, of different domain databases, for a **Paired domain** have the same **Interpro** identifier?

Interpro does not have its own domain models. It defines

Prosite_profiles	222	282	Homeobox domain	PS50071	IPR001356	[Display all genes with this domain]
Smart	224	286	Homeobox domain	SM00389	IPR001356	[Display all genes with this domain]
Pfam	226	281	Homeobox domain	PF00046	IPR001356	[Display all genes with this domain]

domains by the predictions of other domain databases including **Prosite_profiles**, **Smart** and **Pfam**. So if, as here, a **Homeobox domain** is detected by **Prosite_profiles** (**PS50071**), **Smart** (**SM00389**) and **Pfam** (**PF0046**), there exists 3 pieces of evidence to encourage **Interpro** to declare it to believes there to be a **Homeobox domain** (**IPR001356**).

Any one of the **Prosite_profiles**, **Smart** or **Pfam** hits would have been sufficient for **Interpro** to assign membership of this domain to its **Homeobox** classification **IPR001356**.

Where would you expect a **Paired domain** to occur in a protein?

What expectations do you have concerning what typically follows a **Paired domain**?

The **Paired domain** is here said to be found “generally in the N-terminal part” of the protein. That is certainly true of all the examples we have met so far.

The paired domain is an approximately 126 amino acid DNA-binding domain, which is found in eukaryotic transcription regulatory proteins involved in embryogenesis. The domain was originally described as the 'paired box' in the Drosophila protein paired (prd) [(PUBMED:2877747), (PUBMED:3123319)]. The paired domain is generally located in the N-terminal part. An octapeptide [(PUBMED:10811620)] and/or a homeodomain can occur C-terminal to the paired domain, as well as a Pro-Ser-Thr-rich C terminus.

Paired domain proteins can function as transcription repressors or activators. The paired domain contains three subdomains, which show functional differences in DNA-binding. The crystal structures of prd and Pax proteins show that the DNA-bound paired domain is bipartite, consisting of an N-terminal subdomain (PAI or NTD) and a C-terminal subdomain (RED or CTD), connected by a linker. PAI and RED each form a three-helical fold, with the most C-terminal helices comprising a helix-turn-helix (HTH) motif that binds the DNA major groove. In addition, the PAI subdomain encompasses an N-terminal beta-turn and beta-hairpin, also named 'wing', participating in DNA-binding. The linker can bind into the DNA minor groove. Different Pax proteins and their alternatively spliced isoforms use different (sub)domains for DNA-binding to mediate the specificity of sequence recognition [(PUBMED:11103953), (PUBMED:15148315)].

The claim here that “An octapeptide and/or a homeodomain can occur C-terminal to the paired domain, as well as a Pro-Ser-Thr-rich C terminus” confirms what was seen from the **Human PAX** paralogue alignments Previously. That is, sometimes there is a **homeodomain C-terminal** to the **Paired domain**, but not always.

From the **UniProtKB** documentation, you saw that Human **PAX6** at least has “a Pro-Ser-Thr-rich C terminus”

Feature key	Position(s)	Description	Actions	Graphical view	Length
Compositional bias ⁱ	131 – 209	Gln/Gly-rich	Add BLAST		79
Compositional bias ⁱ	279 – 422	Pro/Ser/Thr-rich	Add BLAST		144

Note the mention of the important **prd** Drosophila gene here, overlooked in the **Ensembl** presentation of orthologues to Human **PAX6**?

InterPro did not detect the **Homeobox HTH** as it did the **Paired box HTH**. Have you any thoughts as to why this might be?

The documentation from **SMART**, which really originates from **Interpro**, clearly claims the presence of an **HTH** as the DNA binding element. However, **Interpro** does not predict the presence of an **HTH**, as it did for the **Paired Box**?

I cannot be certain why, however, **HTHs** are difficult to detect just with computer programs. I used to include an exercise that tried for this protein. It proved impossible to obtain a complete picture. One of the reasons being that there are a number of different types of **HTH**. Any given program will typically only search effectively for one type.

Not a very satisfactory answer!

The homeobox domain or homeodomain was first identified in a number of drosophila homeotic and segmentation proteins, but is now known to be well-conserved in many other animals, including vertebrates [(PUBMED:2568852), (PUBMED:1357790)]. Hox genes encode homeodomain-containing transcriptional regulators that operate differential genetic programs along the anterior-posterior axis of animal bodies [(PUBMED:12445403)]. The domain binds DNA through a helix-turn-helix (HTH) structure. The HTH motif is characterised by two alpha-helices, which make intimate contacts with the DNA and are joined by a short turn. The second helix binds to DNA via a number of hydrogen bonds and hydrophobic interactions, which occur between specific side chains and the exposed bases and thymine methyl groups within the major groove of the DNA. The first helix helps to stabilise the structure.

The motif is very similar in sequence and structure in a wide range of DNA-binding proteins (e.g., cro and repressor proteins, homeotic proteins, etc.). One of the principal differences between HTH motifs in these different proteins arises from the stereo-chemical requirement for glycine in the turn which is needed to avoid steric interference of the beta-carbon with the main chain: for cro and repressor proteins the glycine appears to be mandatory, while for many of the homeotic and other DNA-binding proteins the requirement is relaxed.

Why do you suppose there is no match from **PRINTS** or **Prosite** patterns to support the **Homeobox domain** prediction for this protein?

What do you suppose the **Homeobox conserved site** might be?

Contributing signatures

Signatures from InterPro member databases are used to construct an entry.

- **CDD** ⓘ
[cd00086](#) (homeodomain)
- **SMART** ⓘ
[SM00389](#) (HOX)
- **PROSITE profiles** ⓘ
[PS50071](#)
(HOMEBOX_2)
- **Pfam** ⓘ
[PF00046](#) (Homeobox)

Well, the short and rather boring answer to the first part of this question is that **Interpro** did not interrogate **PRINTS** or **Prosite patterns** when it considered the existence of a **Homeobox domain** in this protein!

To prove this you only need to follow the links to the relevant **Interpro** entries and look at the **Contributing signatures**.

Both **PRINTS** and **Prosite patterns** are used to determine the presence of a **Paired domain**. Neither is used to detect a **Homeobox domain**.

Contributing signatures

Signatures from InterPro member databases are used to construct an entry.

- **CDD** ⓘ
[cd00131](#) (PAX)
- **SMART** ⓘ
[SM00351](#) (PAX)
- **PROSITE patterns** ⓘ
[PS00034](#) (PAIRED_1)
- **PROSITE profiles** ⓘ
[PS51057](#) (PAIRED_2)
- **Pfam** ⓘ
[PF00292](#) (PAX)
- **PRINTS** ⓘ
[PR00027](#) (PAIREDBOX)

More interestingly, in the case of **Human PAX6** at least, it would not have made any difference had **PRINTS** and/or **Prosite patterns** been considered for the **Homeobox domain** prediction.

Interpro did actually register a match between the **PAX6 human protein** and the relevant **Prosite pattern**. However, **Interpro** judged this match as too weak (i.e. the probability of a false positive is too high) to be regarded as viable evidence for predicting a **Homeobox domain**. **Interpro** records the match as a conserved site, as you can see from your **Interpro** graphic.

IPR017970 Homeobox, conserved site

Were you to look at the relevant **Prosite** entry (the illustration is a link), you would see that the **Prosite** pattern is quite long, but rather non-specific (the pattern syntax will be fully explained somewhere else). It misses **317** of the **1,639** **Homeobox** domains in **SwissProt**! And

incorrectly claims a **Homeobox** where no **Homeobox** exists on **11** occasions (according to **SwissProt**, which is assumed immaculate in this context). I think **Interpro** is correct to take a hit with this pattern rather lightly.

HOMEBOX_1, PS00027; 'Homeobox' domain signature (PATTERN)

- Consensus pattern:
[LIVMFYGI-ASLVR]-x(2)-[LIVMSTACN]-x-[LIVM]-[Y]-x(2)-[L]-[LIV]-[RKNQESTAIY]-[LIVFSTNKH]-W-[FYVC]-x-[NDQTAH]-x(5)-[RKNAIMW]
- Sequences in UniProtKB/Swiss-Prot known to belong to this class: 1639
 - detected by PS00027: 1322 (true positives)
 - undetected by PS00027: 317 (294 false negatives and 23 'partials')
- Other sequence(s) in UniProtKB/Swiss-Prot detected by PS00027:
11 false positives.

PRINTS has a domain model for **HOMEBOX**, however, it does not match the **PAX6 Human Homeobox domain** sufficiently well to register as a hit! Nearly, but not quite good enough. One might speculate that, in the judgement of **Interpro** at least, the chance of a false **negative** is too high to consider the **PRINTS** model seriously for **Homeobox** detection.

You could just believe me when I claim the **PRINTS HOMEBOX model** does not work in this instance? Instead just speed read the next two pages concentrating only on the last bit which covers the struggles of **PRINTS** to find a **HOMEBOX (recommended)**, or you could prove all for yourself by doing the search. Just for the few doubters and those of you who have nothing better to do, I offer full instructions here (although I feel sure you could work it all out for yourselves).

The **PRINTS** database defines functional protein families. Domains are identified by a number of short, ordered, well-conserved regions. A full match to one of these “fingerprints” will match all the relevant short regions in the correct order. A partial match is recorded if some are missing or if they occur in an incorrect order. **PRINTS** can be searched using the **fingerPRINTscan** program.

Go to the **fingerPRINTscan** home page¹³:

<http://130.88.97.239/PRINTS/>

Select the **FPScan** link and paste in the **PAX6_HUMAN** sequence in raw format. Leave all defaults and hit the **Send Query** button.

Highest scoring fingerprints for your query			
Fingerprint	E-value	GRAPHScan	Motif3D
PAIREDBOX (relations)	1.499643e-43	Graphic	

The top hit is with the **PAIREDBOX** fingerprint. No surprise here. Move down to the list of the best **10** hits.

Ten top scoring fingerprints for your query									
Ancestry	Fingerprint	No. of Motifs	SumId	AveId	PfScore	Pvalue	Evalue	GRAPHScan	
PAIREDBOX	PAIREDBOX	4 of 4	3.5e+02	87	3213	1.3e-49	1.5e-43	IIII	Graphic
HTHREPRESSR	HTHREPRESSR	2 of 2	75.92	37.96	586	5.3e-08	0.17	II	Graphic
POUDOMAIN	POUDOMAIN	2 of 5	65.80	32.90	577	1.7e-07	0.39	...II	Graphic
HOMEobox	HOMEobox	2 of 3	102.06	51.03	724	3e-07	1.2	.II	Graphic
PRICHEXTENS	PRICHEXTENS	3 of 8	102.84	34.28	664	1.2e-05	20	.iIi...	Graphic
POAALLERGEN	POAALLERGEN	2 of 8	42.41	21.20	393	7e-05	1.7e+02i.i	Graphic
TM->GPCRCLAN->GPCRRHODOPSN->LTBRECEPTOR->LTB1RECEPTOR	LTB1RECEPTOR	2 of 6	71.96	35.98	371	0.00032	8.4e+02	..I.I.	Graphic
PROTEINF153	PROTEINF153	2 of 5	52.81	26.40	458	0.00038	6.9e+02	i...i	Graphic
ACONITASE	ACONITASE	2 of 9	63.61	31.80	336	0.00047	1.5e+03iI.	Graphic
GLIADGLUTEN->GLIADIN	GLIADIN	2 of 9	73.82	36.91	396	0.0013	3.7e+03	.I.....I	Graphic

In the list of **Ten top scoring fingerprints**, there is a second **fingerprint** that matches all elements in the correct order. This is the **HTHREPRESSR**. Click on the **HTHREPRESSR** link and from the documentation you can confirm that an **HTHREPRESSOR** is an **HTH** motif of which you might have reasonably expected three? Move back to your **fingerPRINTscan** results. Shimmy down to the **Ten top scoring fingerprints**.



Ten top scoring fingerprints for your query. Detailed by motif									
FingerPrint Name	Motif Number	IdScore	PfScore	Pval	Sequence	Length	low	Pos	high
PAIREDBOX	1 of 4	93.82	815	1.01e-12	VNQLGGVVFVNGRPLPD	16	0	8	0
	2 of 4	82.91	821	6.08e-13	RQKIVELAHSGARPCDISR	19	0	26	0
	3 of 4	87.39	809	2.95e-12	LQVNSGCVSKILGRYYET	18	0	46	0
	4 of 4	83.08	768	6.99e-14	GSIRPRAIGGSKPRVATP	18	0	64	0
HTHREPRESSR	1 of 2	32.91	134	3.98e-02	ARERLAAKID	10	0	239	0
	2 of 2	43.00	452	1.34e-06	DLPEARIQWFSNRRAK	17	0	248	0

From the Position information included in the **Detailed by motif** table, you can see that the **HTH** motif that **fingerPRINTscan** finds is the one that is part of the **HOMEobox** domain it fails to fully detect. **PRINTS** does not see the **HTHs** in the **PAIREDBOX** domain.

¹³ Despite the inexplicitly undecorated URL, this is the currently official **PRINTS** home page provided by **Manchester** University.

Take a look at the **GRAPHScan** for the **PAIREDBOX** prediction and see that it is good! Four out of four very positive motif matches are shown.

Each motif by itself might not be significant. Together, in a **fingerprint**, they constitute a confident prediction for a **Paired Box domain**.



Click on the **Graphic** link for the **HTHREPRESSOR** hit. The best (highest) of the four **motif 1** hits plus the single **motif 2** hit is the finger print that justifies the **HTHREPRESSOR** prediction.

Move back to the **Ten top scoring fingerprints** table. Notice that, whilst there is a prediction for a **HOMEBOX**, it is an incomplete prediction. Only two of the required motifs were detected and so no prediction of a **HOMEBOX** would have been made automatically by **fingerPRINTscan**. This explains why there is no **PRINTS** prediction for a **HOMEBOX** in the **Uniprot Feature Table** for **PAX6_HUMAN**.

However, if you click on the **Graphics** link for the “2 out of 3” motif hit for **homeobox**, you will see that **fingerPRINTscan** only missed the **HOMEBOX** by a whisker! Just a minute bit short for the first motif!

From the Top ten scoring fingerprints table, you can see that **fingerPRINTscan** considers the first motif to be missing (“**.II**”). But I see a fairly healthy **motif 1** in the graphic? I think I would be inclined to give the **HOMEBOX** the benefit of the doubt, would you not? Programs can be so very picky!!! **Its a hit!!**

